

*Multiscale analysis of habitat selection  
by Bonelli's eagle (Aquila fasciata) in NE  
Spain*

**Beatriz Martínez-Miranzo, Eva I. Banda  
& José I. Aguirre**

**European Journal of Wildlife  
Research**

ISSN 1612-4642

Eur J Wildl Res  
DOI 10.1007/s10344-016-1041-x



**Your article is protected by copyright and all rights are held exclusively by Springer-Verlag Berlin Heidelberg. This e-offprint is for personal use only and shall not be self-archived in electronic repositories. If you wish to self-archive your article, please use the accepted manuscript version for posting on your own website. You may further deposit the accepted manuscript version in any repository, provided it is only made publicly available 12 months after official publication or later and provided acknowledgement is given to the original source of publication and a link is inserted to the published article on Springer's website. The link must be accompanied by the following text: "The final publication is available at [link.springer.com](http://link.springer.com)".**

# Multiscale analysis of habitat selection by Bonelli's eagle (*Aquila fasciata*) in NE Spain

Beatriz Martínez-Miranzo<sup>1</sup> · Eva I. Banda<sup>1,2</sup> · José I. Aguirre<sup>1</sup>

Received: 28 January 2016 / Revised: 2 August 2016 / Accepted: 3 August 2016  
© Springer-Verlag Berlin Heidelberg 2016

**Abstract** The habitat selection of 14 individual breeding Bonelli's eagles equipped with satellite tracking devices was evaluated using a multiscale approach during different periods of the annual cycle over 8 years. We studied whether habitat structure and prey availability influence habitat use through the use of vegetation templates and censuses of potential prey. The results showed heterogeneous selection of wooded, rocky, and scrub areas alternating with agricultural areas at a regional scale. At the home range scale, forests and scrubland were mainly selected over the entire year, except during the breeding season, when, surprisingly, humanized areas were selected. Although Bonelli's eagle is considered a forest raptor, during the breeding season they select other types of habitat (i.e., urban areas and dense scrub). This may be related to the high prey availability (especially pigeons) in these areas. Because habitat selection differs at different scales, understanding the effects of this plasticity may be necessary to establish protected areas including urban areas and implement habitat management actions.

**Keywords** GPS satellite telemetry · Multiscale approach · Habitat selection · Compositional analysis · Prey availability

---

✉ Beatriz Martínez-Miranzo  
bmartinez@ucm.es

<sup>1</sup> Department of Zoology and Physical Anthropology Faculty of Biology, Complutense University of Madrid, José Antonio Novais 12, 28040 Madrid, Spain

<sup>2</sup> ENARA Educación Ambiental, XI, 28090 Las Rozas de Madrid, Spain

## Introduction

Spatial and temporal scales in ecology have been included in scientific research for decades (Wiens 1989; Levin 1992). In the field of conservation biology and more specifically in habitat selection studies, the selection of an appropriate scale is very important. Ecological patterns that determine habitat selection may act differently depending on both the spatial scale and temporal scale (Wiens 1989; Levin 1992; Rico Alcázar et al. 2001). Moreover, multiscale approaches may reveal patterns that are not perceived at a single scale (Levin 1992) and may be a determinant in species conservation (Ontiveros et al. 2004).

The use of new tools allows a non-arbitrary scale selection based on biological criteria for the species. The implementation of geographic information systems (GISs), GPS-tracking data, and ecological data has been selected in these types of multiscale habitat selection studies, especially land cover databases (Balbontín 2005). One of the most popular land cover databases in Europe is CORINE. Despite the fact that CORINE is a systematically constructed land cover database covering a large area, it has been shown that this type of land cover data may be insufficient at a detailed scale (Heikkinen et al. 2014). For this reason, it is important to explore particular habitat structures, especially at a local scale where these features may change more rapidly (Wiens 1989). In addition, comparisons should be made with the available digital land cover information.

Similar to habitat structure, climate and resource availability can influence habitat selection as well (Ontiveros and Pleguezuelos 2000; Ontiveros et al. 2005; López-López et al. 2006). Territorial species establish their home range based on resource availability, for example the availability of nesting areas (López-López et al. 2006) and prey

(Ontiveros and Pleguezuelos 2000). However, this resource availability may vary over the years or over particular periods in a single season. Recording food availability and its distribution throughout the home range can help to understand not only occurrence patterns of individuals at a particular place (regional scale) or the establishment of their territories (home range scale) but also their particular use of resources within the home range (local scale).

This is the case of Bonelli's eagle (*Aquila fasciata*), a territorial raptor that is distributed throughout the western Palearctic but mainly restricted to the Mediterranean region (Hagemaijer and Blair 1997; Ontiveros 2014). In the last several years, it has suffered a general decline in its populations (BirdLife International 2016), but most severely in the western area of the Iberian Peninsula (Ontiveros 2014). Changes in land use by humans and a decrease in potential prey availability have played an important role in their decline (Ontiveros 2014).

Studies about habitat selection by Bonelli's eagle are a key to gaining knowledge about the spatial ecology of this species. Muñoz et al. (2005) and Carrascal and Seoane (2009) indicated the factors affecting the distribution of this species at a large scale using geographic, climatic, landscape, and human variables. On the other hand, Carrete et al. (2002) and López-López et al. (2006) explored habitat preference factors at a local scale also using these types of variables. Balbontín (2005) used the same approach to study juvenile dispersal. To our knowledge, this is the first study that uses precisely defined home ranges (Martínez-Miranzo et al. 2016) of 14 adult individuals of different sexes at different spatial and temporal scales.

The aim of this study is to evaluate habitat selection by Bonelli's eagle at different spatial and temporal scales and whether factors like habitat structure and prey availability determine long-term habitat selection. In accordance with that, the results of this study may have important repercussions in the knowledge about the spatial ecology of this eagle, helping to establish appropriate conservation policies.

## Methods

### Study area

The study was conducted in the Aragon Region, Northeast Spain. The altitude in the area ranges from 130 to 1200 m above sea level. Land cover consists mainly of coniferous forests and large areas of Mediterranean scrub filled with cultivation areas, mostly of dry cereals, fruit trees, and Mediterranean crops (olive trees and vineyards). Craggs, cliffs,

and other unproductive areas like steppes are also present in this area (Sampietro et al. 1998).

### Data collection

From 2004 to 2013, 14 adult breeders of Bonelli's eagles (eight males and six females) were trapped in Aragon using radio-controlled bow-net traps. All individuals were ringed with a metal ring and were equipped with a 45-g Argos/GPS PTT device (Microwave Telemetry, MD, USA). Transmitters were powered with solar panels and fixed to birds as backpacks with a Teflon harness with a central ventral rupture point (Garcelon 1985). The weight of the transmitters only represents 2.25 % of the total body weight (Kenward 2001). PTTs were programmed to work between 0600 hours and 2100 hours, and collect one location per hour. To avoid bias towards roosting areas, consecutively repeated locations in the early morning and late evening of inactive eagles were excluded because they were considered to be non-independent (Swihard and Slade 1985; Seaman and Powell 1996; Kenward 2001). A total number of 59,482 locations from the 14 individuals were obtained.

### Multiscale and temporal habitat selection

The size and shape of the home range between years is maintained by Bonelli's eagles in this area, but there are variations in the use within the home range depending on the period of year (Martínez-Miranzo et al. 2016). The analysis of habitat selection was conducted at three different temporal scales and spatial levels of detail according to Johnson (1980) (regional scale, including all Aragon geographical regions; study area scale, including all spaces with a valid location obtained by GPS; and home range scale, within each territory calculating from GPS data; hereafter, RS, SA, and HR, respectively).

For the temporal variations in habitat selection, we divided the year into three periods related to the biological cycle of the species (Arroyo et al. 1995). Period 1 was defined as the non-breeding season (NBr), from September 1 to February 14, when breeding individuals are less tied to their nesting area and made distant movements (Ontiveros 2014). In period 2 or the breeding season (Br) (from February 15 to June 14), both parents invest in clutches but females spend most of the time at the nest, and in general, parents' movements are restricted (Ontiveros 2014). During period 3 or post-fledging dependence period (Pfd), from June 15 to August 31, parents continue to feed fledglings near nesting areas until the juveniles leave the territories where they were born and disperse (Real et al. 1998).

The different habitat types were extracted following habitat structure criteria from previous Bonelli's eagle preferences (Ontiveros 2014) from categories in CORINE Land Cover (CLC) (European Environment Agency 2007) depending on the scale used for the analysis (CLC 2006 for regional and



study area scales and CLC 2000 for home range scale). We were unable to use the same CLC data for all the analysis because the detail level of CLC 2006 is lower than later versions of CLC 2000 (Table 1). In order to establish more precise habitat structure preferences at a home range scale, the three highly selected categories for study area scales (forest, scrub, and grassland) were redefined more precisely into nine new categories following CLC 2000 (i.e., study area: scrub was redefined at a home range scale into dense scrub, open scrub, coniferous scrub, and hardwood scrub) (Table 1). The number of categories were restricted according to data analysis used (Aebischer et al. 1993).

To test for random habitat selection by breeders at a RS, we performed a chi-square analysis in Statistica 8.0 software (StatSoft Inc. 2007). Using random point generation in ArcGIS 9.3 software (ESRI 1999–2009), we generated

the same number of random points as GPS locations in all Aragon regional areas and tested the frequency difference between the two data sets. ANOVA analysis in Statistica 8.0 software was selected to test the temporal variation at this scale.

To perform habitat selection analysis at the SA level, we built a minimum convex polygon (MCP, 100 %) defined as the maximum area used by individuals (Kenward 2001). MCP was calculated with all valid locations including outermost locations. Individual home range was estimated using Hawth's tools (Beyer 2004) and fixed kernel methods, 95 % isopleths (Worton 1989) with a default smoothing factor of 1 (Fernández et al. 2009; Bosch et al. 2009; Martínez-Miranzo et al. 2016) in ArcGIS 9.3 software. Home range sizes were constructed using only diurnal locations.

**Table 1** Habitat type composition and percentage of different habitat categories extracted from CLC depending on the scales CLC 2006 (regional scale and study area scale) and CLC 2000 (home range scale)

Habitat type composition	CLC 2006	Percent	CLC 2000	Percent
Scrub	Transitional woodland shrub	3.67	Coniferous scrub	5.13
	Sclerophyllous vegetation	12.15	Dense scrub	19.32
			Open scrub	35.62
			Mixed scrub	0.07
Forest	Broad-leaved forest	2.47	Hardwood scrub	1.04
	Coniferous forest	7.44	Evergreen forest	3.19
	Mixed forest	0.29	Coniferous forest	33.65
			Riparian forest	0.74
Grassland	Natural grassland	0.99	Natural grassland	1.24
Crops	Non-irrigated arable land	40.97	Non-considered	
	Permanently irrigated land	9.34	Non-considered	
	Rice fields	0.52	Non-considered	
	Annual crops	0.00	Non-considered	
	Complex cultivation	7.63	Non-considered	
	Crops and natural vegetation	8.01	Non-considered	
Fruit crops	Vineyards	1.37	Non-considered	
	Fruit trees	1.05	Non-considered	
	Olive groves	1.09	Non-considered	
Urban	Urban continuous	0.31	Non-considered	
	Urban discontinuous	0.14	Non-considered	
	Industrial area	0.25	Non-considered	
	Human networks	0.04	Non-considered	
	Mineral extraction	0.12	Non-considered	
	Dump sites	0.02	Non-considered	
	Construction sites	0.15	Non-considered	
	Ocio area	0.02	Non-considered	
Water	Inland waters	0.29	Non-considered	
	Water bodies	0.37	Non-considered	
Bare rock	Bare rock	0.09	Non-considered	
Unproductive	Sparsely vegetated areas	1.13	Non-considered	
	Burnt areas	0.03	Non-considered	
	Dunes and sand plains	0.05	Non-considered	

Compositional analysis described by Aebischer et al. (1993) was selected to study habitat selection at SA and HR levels. This analysis utilizes a MANOVA test to compare the proportion of habitat available for use and shows a rank of habitat types in order of use. In the cases where the habitat value is zero (not available or no use), we used the value 0.01 as recommended in Aebischer et al. (1993).

We conducted vegetation templates within the study area to find differences in habitat structure at an HR scale between CLC 2000 and actual composition. Following the method described by Prodon and Lebreton (1981), we recorded the vegetation structure along 140 randomly selected transect (2.5 km approx. each). In total, 1033 vegetation templates were made at the beginning and end of each itinerary and each time there was contact with any potential prey. Line transects were performed on foot during two consecutive years in the three annual periods previously described. We visually estimated grass cover (the percent of vegetation below 0.5 m in height), scrub cover (the percent of vegetation between 0.5 and 2 m in height), and tree cover (the percent of vegetation above 2 m in height).

Only scrub cover was selected for the analysis because scrubland has a positive effect on the frequency of species occurrence (Carrascal and Seoane 2009) and is one of the most selected habitat types at this scale. With the percent of vegetation structure calculated in each transect, we created two categories in relation to the principal type of scrub cover in CLC 2000. Values between 0 and 40 % were selected because they best fit the values recorded by CLC 2000. Open scrub was assigned to percent between 0 and 40 % and dense scrub to percent between 40 and 100 %. We compared whether there were differences between scrub cover in CLC 2000 and the actual scrub cover. In addition, we checked for the possible difference between periods and years.

### Prey availability

To record prey availability at different habitat types, we selected the main prey groups for this species in Aragon. Pigeons (including *Columba* sp. and *Streptopelia* sp.) (27 %), lagomorphs (including *Oryctolagus cuniculus* and *Lepus europaeus*) (22 %), partridges (*Alectoris rufa*) (11 %), and corvids (*Corvus* sp.) (7 %) (Alcántara et al. 2003) represent up to 67 % of Bonelli's eagle diet in Aragon. Direct censuses on foot were performed (Tellería 1986). A total of 140 random transect (2.5 km approx. each) were performed during two consecutive years in the three annual periods described above in the study area. A total of 1050 km were censused, and 753 contacts of prey were obtained. The very low presence of rabbit and partridge in the study area was insufficient for analysis. For each itinerary, the total number of available prey was recorded and corrected by the total

length of each transect obtaining an index of prey/length unit (kilometric abundance index, KAI) (Tellería 1986).

We compared prey availability with scrub habitat type. This type of habitat may influence the presence and detectability of prey by the eagles. To overcome the large number of no prey presence in the transect, generalized linear model (GLM) analysis in Statistica 8.3 software with Poisson distribution and logit transformation was performed. Prey type was used as a dependent variable, and the presence of clear and dense scrub was the categorical explanatory variable. For all statistical tests, probability values less than 0.05 were considered significant.

## Results

### Habitat selection

At the regional scale, habitat selection by Bonelli's eagle showed a strong tendency towards scrub and forest, which represent 76.5 % of the total habitat selection. Results differed significantly from random ( $\chi^2_3 = 68874.42$ ,  $p < 0.001$ ). No differences between periods were found at this scale.

Compositional analysis at the study area scale showed that eagles do not use the habitat randomly. We found significant differences in habitat use among the three periods of the year (see Table 2). According to the ranking matrix, forest and scrub habitat were the most used while agricultural areas like fruit trees and crops were less selected. Nevertheless, we detected differences in selection order between periods (Table 2). Forest was selected more than scrub outside of the breeding season, while during the breeding season, scrub and rock were the most chosen habitats. In addition, urban areas were significantly more preferred during the breeding season.

We also found significant values at a home range scale (Table 2). Compositional analysis showed that coniferous forest and dense scrub were the most selected and evergreen and riparian forests were the least preferred habitats. Differences in use between periods were also found. Dense scrub is more selected during the breeding season and post-fledging dependence period, while coniferous forest was the most preferred during the non-breeding season.

We found significant differences between scrub cover in different periods ( $F_{(2,631)} = 7.6649$ ;  $p < 0.001$ ). The scrub cover values were higher during the breeding season and lower during the non-breeding season. No differences were found between actual scrub cover categories and CORINE categories ( $F_{(1,631)} = 0.00063$ ;  $p = 0.979$ ). The scrub cover values did not change between years.

**Table 2** Ranked matrix of habitat type selection for all individuals ( $n = 14$ )

Scale	Period	Wilk's $\lambda$	$p$	Ranked habitat types
SA	NBr	0.1046	0.0090	FOR > SCR > ROC > GRA > WAT > URB > UNP > FRU > CRO
	Br	0.0937	0.0020	SCR > ROC > FOR > URB > GRA > WAT > UNP > FRU > CRO
	Pfd	0.0547	0.0010	FOR > SCR > WAT > ROC > GRA > URB > UNP > FRU > CRO
HR	NBr	0.3324	0.0355	CON.F > CON.S > GRA > DEN.S > OPE.S > MIX.S > HAR.S > RIP.F > EVE.F
	Br	0.2893	0.0171	DEN.S > GRA > CON.F > MIX.S > CON.S > OPE.S > RIP.F > EVE.F > HAR.S
	Pfd	0.2857	0.0160	DEN.S > MIX.S > CON.S > GRA > CON.F > OPE.S > HAR.S > RIP.F > EVE.F

For study area scale (SA), the habitat types are forest (FOR), scrub (SCR), bare rock (ROC), grassland (GRA), water (WAT), urban (URB), unproductive (UNP), crop fruit (FRU), and crops (CRO), and for home range scale (HR), the habitat types are coniferous forest (CON.F), coniferous scrub (CON.S), grassland (GRA), dense scrub (DEN.S), open scrub (OPE.S), mixed scrub (MIX.S), hardwood scrub (HAR.S), riparian forest (RIP.F), and evergreen forest (EVE.F)

### Prey availability

GLZs showed significant differences between pigeon abundance and habitat structure. Higher abundances of pigeons were found in dense scrub (Wald  $\chi^2_1 = 17.563$ ,  $p < 0.001$ ). On the other hand, when we compared corvid abundance and habitat structure, they showed higher abundances in clear scrub (Wald  $\chi^2_1 = 5.6962$ ,  $p = 0.017$ ).

### Discussion

This study shows the importance of a multiscale approach to identify habitat selection by Bonelli's eagle. Our results show that while, at a regional scale, individuals select heterogeneous habitat with crops areas, scrub areas, and coniferous forest, at a smaller scale, habitat structure within the home range plays a key role in habitat selection. The increased use of scrubland and coniferous forest, as with other areas with human presence, has been detected. Selection seems to be conditioned by the presence of potential prey and personal experience of each individual. Such selection varies depending on the season and the needs of individuals at each particular moment of the season.

The integration of modern tracking tools and classical census methods provides large amounts of high-quality data. This allowed us to implement the method described by Aebischer et al. (1993), avoiding its main problems (i.e., inappropriate level of sampling and sample size, non-independence of proportions, and arbitrary definition of habitat availability). It also allowed us to establish sampling periods synchronized with the biological cycle of the species.

Similarly, studies involving comparisons over time can reveal differences in habitat use related to the needs of each species at a particular time during the annual cycle (e.g., breeding season in raptors). For this reason, it is important to consider

seasonal variability in the use of space and should be linked to the availability of resources and the importance of a heterogeneous and changing habitat within a study area. Therefore, long-term studies of endangered species are also important because conservation policy implementation in large areas is often based on very short-term studies (Wiens 1989).

At a regional scale, we found a non-random selection of habitat types. In line with other studies (Carrascal and Seoane 2009; Ontiveros 2014), Bonelli's eagle in the Aragon region selected heterogeneous landscapes with scrub and forest, dotted with cliffs (important for nest site selection by this raptor) (López-López et al. 2003). Prey detectability seems to be the main factor driving the selection of this type of habitat (Ontiveros et al. 2005). Nevertheless, crops and other fruit fields were not selected by individuals (Carrete et al. 2002). Despite the fact that this species can tolerate human presence (Muñoz et al. 2005), high-intensity human activities such as agricultural practices or heavy vehicle traffic in the area may exceed the eagles' tolerance threshold, regardless of higher prey abundance (pigeons, partridges, and rabbits in fruit crops and edge habitats) (authors' unpublished data). Furthermore, no temporal variation was found at this scale. This variation is difficult to detect at a large scale and even at other levels.

At the study area scale, eagles showed a differing habitat selection among seasonal periods. Rocks were selected by individuals during the breeding season. Bonelli's eagle is a Mediterranean raptor that nests in cliffs at moderate altitudes, and therefore, a positive selection for this habitat is expected during this period. Scrub was also more selected during this period. The presence of chicks during the breeding season demands provision of high amounts of food by the breeders. Scrub is the preferred habitat for the main prey species of Bonelli's eagle (rabbits and partridges) (Gil-Sánchez et al. 2000; Carrete et al. 2002). Therefore, individuals spend more time in these areas hunting. Forests (principally coniferous forests) are more selected during the rest of the periods.

Although they do not visit the nest area frequently, they spend a lot of time in forest habitat during the rest of the year, primarily for roosting and defending their home range.

Urban areas (small rural villages and open industrial areas) were primarily selected during the breeding season over other habitats. The scarce abundance of prey for these eagles (rabbits and partridges) in their original habitats and the plasticity of this species to adjust their diet can condition such selection (Ontiveros and Pleguezuelos 2000). Under conditions of prey shortage, Bonelli's eagles can hunt rock pigeons (*Columba livia*) and common woodpigeons (*Columba palumbus*). Pigeons concentrate mainly in urban habitats (Palma et al. 2006), and therefore, eagles use these high-density areas to hunt more efficiently. In fact, there is an important percent of this type of prey in the diet of Bonelli's eagle in Aragon (Alcántara et al. 2003).

Individuals' experience, especially in raptors with large home ranges, is important to optimize resource exploitation. At the home range scale, we found that dense scrub is more selected than open scrub. In contrast to other studies (Balbontín 2005; López-López et al. 2006), breeders in Aragon preferred this type of scrub although prey detectability is lower. In spite of the fact that the main prey such as rabbits and partridges is very common in areas with clear Mediterranean scrub, alternative prey such as pigeons (which makes up 26.7 % of the diet in Aragon (Alcántara et al. 2003)) is also associated with coniferous forest and transition areas with dense scrub. The shortage of main prey in the study area along with the personal experience of the individuals and the knowledge of their home range can lead individuals to spend more time looking for alternative prey such as pigeons in these areas of dense scrub despite their lower detectability.

In conclusion, long-term multiscale habitat selection studies can reveal aspects that are undetected at a single scale or that might need some time to be revealed due to changes during the year mainly driven by differential resource availability. In addition, the use of new tracking technology can show more precise results in certain areas and can address more precise conservation concerns. In our study area, we confirmed that in spite of the fact that individuals follow a general pattern for establishing home range, prey availability is very important to determine that home range. The home range use by individuals is closely related to the period of the year. Therefore, it is very important to implement conservation measures not only at a large scale but also at a short time scale, keeping in mind the variation throughout the year. Habitat structure and the adaptation of the species to habitat changes should be considered. For example, the use of urban areas by Bonelli's eagles during the breeding season is not usually included in conservation programs. In the same way, conservation policies addressing temporal variation could be considered, for example regulating climbing activities during the breeding season and managing forest areas during the non-breeding season.

**Acknowledgments** We would like to thank the Government of Aragon for providing all study data, especially M. Alcántara and D. Guzman. Our thanks to all Nature Protection officers, APNs, involved in the eagles' management in Aragon. We want to extend our special thanks to E. Ferreiro y A. Gardiazabal, a BIOMA TBC consultant, for all the support during the study and outside of it.

## References

- Aebischer NJ, Robertson PA, Kenward RE (1993) Compositional analysis of habitat use from animal radio-tracking data. *Ecology* 74:313–325
- Alcántara M, Ferreiro E, Gardiazabal A (2003) El Águila-azor Perdicera en Aragón. *Naturaleza Aragonesa* 10:41–47
- Arroyo B, Ferreiro E, Garza V (1995) El águila perdicera *Hieraetus fasciatus* en España. Censo, reproducción y conservación. Ministerio de Agricultura Pesca y Alimentación. Colección Técnica: ICONA
- Balbontín J (2005) Identifying suitable habitat for dispersal in Bonelli's eagle: an important issue in halting its decline in Europe. *Biol Conserv* 126:74–83
- Beyer HL (2004) Hawth's analysis tools for ArcGIS. (Accessed online [www.spatialecology.com/htools](http://www.spatialecology.com/htools))
- BirdLife International (2016) Species factsheet: *Aquila fasciata*. Downloaded from [www.birdlife.org](http://www.birdlife.org). Accessed 9 Aug 2016
- Bosch R, Real J, Tinto A, Zozaya EL, Castell C (2009) Home-ranges and patterns of spatial use in territorial Bonelli's eagles *Aquila fasciata*. *Ibis* 152:105–117
- Carrascal LM, Seoane J (2009) Factors affecting large-scale distribution of the Bonelli's eagle (*Aquila fasciata*) in Spain. *Ecol Res* 24:565–573
- Carrete M, Sánchez-Zapata JA, Martínez JE, Sánchez MA, Calvo JF (2002) Factors influencing the decline of a Bonelli's eagle (*Hieraetus fasciatus*) population in southeastern Spain: demography, habitat or competition? *Biodivers Conserv* 11:975–985
- European Environment Agency (2007) CLC2006 technical guidelines., p 66
- Fernández M, Oria J, Sánchez R, González LM, Margalida A (2009) Space use of adult Spanish imperial eagles *Aquila adalberti*. *Acta Ornithol* 44:17–27
- Garcelon DK (1985) Mounting backpack telemetry packages on bald eagles. Institute for Wildlife Studies, Arcata
- Gil-Sánchez JM, Molino-Garrido F, Valenzuela-Serrano G, Moleón M (2000) Demografía y alimentación del Águila-Azor Perdicera (*Hieraetus fasciatus*) en la provincia de Granada. *Ardeola* 47:69–75
- Hagemajier EJM, Blair MJ (1997) The EBCC atlas of European breeding birds: their distribution and abundance. T & A D Poyser, London
- Heikkinen RK, Bocedi G, Kuussaari M, Heliölä J, Leikola N (2014) Impacts of land cover data selection and trait parameterisation on dynamic modelling of species' range expansion. *PLoS One* 9(9):e108436
- Johnson D (1980) The comparison of usage and availability measurements for evaluating resource preference. *Ecology* 61(1):65–71
- Kenward RE (2001) A manual for wildlife radio tagging. Academic, London
- Levin SA (1992) The problem of pattern and scale in ecology. *Ecology* 73:1943–1967
- López-López P, García-Ripollés C, Aguilar J, García-López F, Verdejo J (2003) Two-spatial scale multifactor analysis on Bonelli's eagle (*Hieraetus fasciatus*) nest-site selection in the east of Iberian Peninsula. In: Proceedings II Iberian ornithological workshop. Aveiro, Portugal
- López-López P, García-Ripollés C, Aguilar J, García-López F, Verdejo J (2006) Modelling breeding habitat preferences of Bonelli's eagle (*Hieraetus fasciatus*) in relation to topography, disturbance, climate and land use at different spatial scales. *J Ornithol* 147:97–106



- Martínez-Miranzo B, Banda E, Gardiazábal A, Ferreiro E, Aguirre JI (2016) Differential spatial use and spatial fidelity by breeders in Bonelli's eagle (*Aquila fasciata*). *J Ornithol* (in press)
- Muñoz AR, Real R, Barbosa AM, Vargas JM (2005) Modelling the distribution of Bonelli's eagle in Spain: implications for conservation planning. *Divers Distrib* 11:477–486
- Ontiveros D (2014) Águila perdicera—*Hieraaetus fasciatus*. In: Salvador A, Morales MB (eds) Enciclopedia Virtual de los Vertebrados Españoles. Museo Nacional de Ciencias Naturales, Madrid, Accessed online [www.vertebradosibericos.org](http://www.vertebradosibericos.org)
- Ontiveros D, Pleguezuelos JM (2000) Influence of prey densities in the distribution and breeding success of Bonelli's eagle (*Hieraaetus fasciatus*): management implications. *Biol Conserv* 93:19–25
- Ontiveros D, Real J, Balbontín J, Carrete MR, Ferreiro E, Ferrer M, Mañosa S, Pleguezuelos JM, Sánchez-Zapata JA (2004) Conservation biology of the Bonelli's eagle in Spain: research and management. *Ardeola* 51:461–470
- Ontiveros D, Pleguezuelos JM, Caro J (2005) Prey density, prey detectability and food habits: the case of Bonelli's eagle and the conservation measures. *Biol Conserv* 123:19–25
- Palma L, Beja P, Pais M, Da Fonseca LC (2006) Why do raptors take domestic prey? The case of Bonelli's eagles and pigeons. *J Appl Ecol* 43(6):1075–1086
- Prodon R, Lebreton JD (1981) Breeding Avifauna Mediterranean succession: the holm oak and cork oak series in the eastern Pyrenees. 1. Analysis and modelling of the structure gradient. *Oikos* 37:21–38
- Real J, Mañosa S, Codina J (1998) Post-nestling dependence period in the Bonelli's eagle (*Hieraaetus fasciatus*). *Omis Fennica* 75:129–137
- Rico Alcázar L, Martínez JA, Morán S, Navarro JR, Rico D (2001) Preferencias de hábitat del Águila-azor Perdicera (*Hieraaetus fasciatus*) en Alicante (E de España) a dos escalas espaciales. *Ardeola* 48:55–62
- Sampietro FJ, Pelayo E, Hernández F, Cabrera M, Guiral J (1998) Aves de Aragón. Atlas de especies nidificantes. Diputación General de Aragón, Ibercaja, Zaragoza, p 588
- Seaman DE, Powell RA (1996) An evaluation of the accuracy of kernel density estimators for home range analysis. *Ecology* 77:2075–2085
- StatSoft Inc. (2007) STATISTICA (data analysis software system), version 8.0., [www.statsoft.com](http://www.statsoft.com)
- Swihard RK, Slade NA (1985) Testing for independence of observations in animal movements. *Ecology* 66:1176–1184
- Tellería JL (1986) Manual para el censo de los vertebrados terrestres. Raíces, Madrid
- Wiens JA (1989) Spatial scaling in ecology. *Funct Ecol* 3:385–397
- Worton BJ (1989) Kernel methods for estimating the utilization distribution in home-range studies. *Ecology* 70:164–168