

Abstracts from CIP 2009

Tercer Congreso Ibérico de Percepción

SYMPOSIUM 1: BIOLOGICAL MOTION

Moderator: Jorge Santos

Recognition of Body Motion: From Visual Sequence Analysis to Structured Dynamic Motor Representations.

M.A. Giese. (Department of Cognitive Neurology, Hertie Institute for Clinical Brain Research & CIN, Tübingen, Germany).

The recognition of biological motion and actions integrates processes at multiple cortical levels. On the one hand, and consistent with physiologically inspired neural models, accumulating evidence suggests a critical influence of the visual processing of temporal sequences of form and motion patterns. On the other hand, a very extensive literature has emerged showing influences of motor representations on the visual recognition of imitable actions. The exact relationship between these different representations remains largely unknown. We will first show how learning-based visual processing mechanisms account not only for the processing of biological-motion stimuli, but also for the recognition of goal-directed actions, at the same time reproducing critical visual tuning properties of “mirror neurons” in premotor cortex. We will then show that biological motion recognition is influenced by concurrently executed motor behavior in a spatio-temporally highly selective manner. This indicates the existence of highly structured interactions between dynamic representations for the recognition and execution of motor behavior. We finally will show how constraints, potentially resulting from the motor control of complex full-body movements, can be exploited for a detailed study of critical features that determine the perception of emotions from body movement.

Multimodal perception and action in biological motion

J.A. Santos (Department of Psychology, University of Minho, Portugal), S. Mouta (Department of Psychology, University of Minho, Portugal), C. Mendonça (Department of Psychology, University of Minho, Portugal) and J. Ferreira (Department of Industrial Electronics, University of Minho, Portugal).

Most of the studies on biological motion perception are concerned with recognition issues and visual-only inputs. However, perceptual cues from several modalities and action-related tasks are critical and common in everyday life situations. Here, we present two research approaches, one on multimodal perception (visual and auditory) and the second on the role of observers' movement. In an audiovisual biological motion experiment, we found that sound strongly affects our final multimodal direction judgments. We used point-light-walkers with a strong bias in order to be perceived as walking towards the observer. We then combined these stimuli with congruent looming or receding auditory steps. In the audiovisual condition, we found that the bias was eliminated. In a time-to-passage (TTP) estimation experiment, a self-based TTP judgment between an external visual object and the observer was studied in static and walking conditions. Walking observers were more accurate and prompt to estimate time-to-passage when compared with static ones. In summary, both multimodal cues and action seem to play a relevant role in achieving accurate biological motion perception.

Translational biological motion stimuli for perceptual research

B. Aragão (Universidade do Minho, Instituto de Educação e Psicologia), L. Fontes (Universidade do Minho, Instituto de Educação e Psicologia), R. Pereira (Universidade do Minho, Instituto de Educação e Psicologia), J.A. Santos (Universidade do Minho, Instituto de Educação e Psicologia), E. Soares (Universidade do Porto, Faculdade de Engenharia) and M.V. Correia (Universidade do Porto, Faculdade de Engenharia and Instituto de Engenharia de Sistemas e Computadores).

Since the studies of Johanson in 1973, the standard stimulus for biological motion perception has been the Point-Light Walker (PLW). This consists of a pattern of moving dots with the common translational component removed as if the target person were walking on a treadmill. The translational motion component has remained as a matter of interest only for researchers on biomechanics and its applications in fields such as rehabilitation, sports training, or virtual actor designs. Nevertheless, there are specific features of the translational path that may be useful for our visual system in several tasks such as recognition, motion perception, and action control. In this communication, we will highlight relevant visual features of the translational component in biological motion and we will address motion capture conceptual and methodological issues.

Neural correlates of 2D and 3D biological versus non-biological object perception: Psychophysics and neurophysiology

B. Graewe (IBILI, Faculty of Medicine, Coimbra, Portugal and Department of Cognitive Neuroscience, Faculty of Psychology, University of Maastricht, The Netherlands), R. Farivar (McGill University, Canada), P. de Weerd

(Department of Cognitive Neuroscience, Faculty of Psychology, University of Maastricht, The Netherlands) and M. Castelo-Branco (IBILI, Faculty of Medicine, Coimbra, Portugal).

Event-related potential studies can provide important insights into the understanding of how and when the human brain differentiates between visual stimulus categories. Here, we ask the question whether ERP components related to object recognition show category-specific responses. Many studies have concluded that faces elicit such category-specific responses, based on the demonstration of a “face-specific” negative ERP potential termed the “N170,” which is more pronounced for faces than for other visual stimuli. In the present study (8 participants), using static stimuli (photographs of faces and chairs and Mooney faces) and manipulating their saliency by different levels of masking, we found that the amplitude of the N170 could be modulated by the saliency of the stimuli. Hence, whether the N170 of faces was larger than for non-faces simply depended on the saliency level of the stimuli. Additionally, using structure-from-motion (SFM) defined face and chair stimuli, we found an object-related negative peak at 250 ms whose amplitude could be modulated by varying depth of the respective stimulus categories (“flattening” the stimuli leading to increasingly smaller amplitudes). Moreover, a comparison between stimulus categories (faces, chairs, and coherent motion) revealed that the N250 peak has a significantly larger amplitude for SFM chair stimuli than for SFM face stimuli (even for normal/optimized depth values in the two categories) and hence failed to show a face-specific effect. The object-sensitive N250 elicited by the SFM stimuli peaked around 80 ms later compared to the N170 in the static face condition, which can be explained by a perceptual delay in perceiving SFM stimuli (i.e., the time needed in order to detect an object from the moving dot pattern). We therefore speculate that the N250 we found for SFM stimuli is a perceptual analogue of the N170 for static faces. Support for this claim comes from our finding that the N170 for static faces/chairs can also be shifted depending on the exact time of their presentation (in an experimental condition in which static stimuli become detectable later in the stimulus period, simulating the SFM condition). Taken together, these findings demonstrate a strong dependence of object-related ERP signals on depth/saliency of the respective stimulus category and a lack of an unambiguous relationship between the amplitude of object-sensitive ERP components and object categories.

SYMPOSIUM 2: VISUAL ILLUSIONS AND COMPLEXITY

Moderator: Enric Munar

Illusions and illusionists: How to fool the brain with magic and other tricks

S. Martínez-Conde (Barrow Neurological Institute, Phoenix, United States).

All our life, every object we see, every person we know, and every incident we experience are just the product of our imagination. Each and every one of our thoughts and feelings are the result of brain processes, and not necessarily the result of an event in the real world. The same neural machinery that interprets the sensory inputs also creates our thoughts, imaginations, and dreams; thus, the world we experience and the world we imagine have the same physical bases in the brain. Just as physicists study the most minute subatomic particles and the largest galactic conglomerates to understand the universe, neuroscientists must examine the cerebral processes underlying perception to understand our experience of the universe. Visual illusions are one of our most important tools to understand how the brain builds our experience of reality. Likewise, the principles developed by magicians and illusionists throughout history can be very useful to manipulate attention and awareness in the laboratory. I will discuss how the visual and cognitive illusions developed by artists and magicians can be applied to the study of the neural bases of consciousness and perception.

Visual complexity and beauty

M. Nadal and E. Munar (Human Evolution and Cognition. Associated group to IFISC (UIB-CSIC)).

The influence of complexity on aesthetic preference and the appreciation of beauty has been noted since Classical Antiquity. However, the scientific formulation of this influence dates to the late nineteenth and early twentieth centuries, with the birth of empirical aesthetics. Although the influence of complexity on the appreciation of beauty became pivotal in Daniel Berlyne’s *Psychobiological Aesthetics* framework, empirical studies attempting to determine the nature of this relation have produced divergent results. Whereas in some instances, the inverted-U function posited by Berlyne was found, in many other cases, an ascending or descending linear relation resulted from the experiments. We will argue here that this discrepancy is due to differences in the way the concept of complexity has been defined and operationalized. Based on the notion that there are several different aspects of complexity, one related with the amount of elements in the stimulus, another with the organization of those elements, and another with symmetry, we obtained curve fit measurements for ratings of 60 images on a beauty

scale and three complexity factors (amount of elements, organization, and symmetry). Results showed that these different forms of complexity had different effects on the appreciation of beauty. Whereas rated beauty increases linearly with the amount of elements, it had a U-shaped relation with organization, and an inverted-U relation with symmetry. We conclude from these results that a general concept, one that does not distinguish among these three aspects, is unsuited for studying the effects of visual complexity on people's appreciation of beauty.

Adjustment strategies in the Vertical-Horizontal illusion

J.E. García-Albea, D. Rivera, J. M. Gavilán, M. Guasch and J. Demest (Dept. of Psychology/CRAMC - Universitat Rovira i Virgili (Tarragona, Spain)).

One of the most persistently studied perceptual illusions is the Vertical-Horizontal (VH) illusion, according to which people tend to overestimate the length of a vertical line when compared with the length of a horizontal line of exactly the same magnitude. The most frequent procedure used to test the presence of the illusion and to assess its size is based on the psychophysical *method of adjustment*, by means of the resulting average error when trying to equate the length of a variable line (V or H) to the length of a standard one (H or V, respectively). The work presented here makes use of this method in order to examine the relative weight of different factors that may contribute to the illusory effect, such as (a) the frame of reference surrounding the stimulus lines, (b) the proper direction of the length adjustment demanded by the task (V > H or H > V), and (c) the presence (or not) of supervised practice through feedback. Furthermore, the illusory effect has been tested and measured across four different VH configurations: (a) a cross, (b) an L, (c) an inverted T, and (d) a T rotated counterclockwise 90°. Our data show that the size of the expected illusion increases from Figure 1 to Figure 2, and from the latter to Figure 3, where it reaches its maximum; but in Figure 4, the illusion reverses, reaching a significant effect in the opposite direction. This pattern of results appears consistently across different stimulus and test conditions, even though the variability in the amount of illusory effect seems to be modulated by the intervention of the critical factors just examined. Their relevance for a better understanding of the mechanisms responsible for the illusion will be discussed, with a look at the modularist characterization of the early visual processes.

Supported by AGAUR, SGR2005-471 (Generalitat de Catalunya)

What drives the first dominance phase in binocular rivalry?

M. Blanco (Departamento de Psicología, Universidad de Santiago de Compostela, Spain), F. Valle-Inclán (Departamento de Psicología, Universidad de La Coruña, Spain), J. Corral (Departamento de Psicología, Universidad de La Laguna, Tenerife, Spain) and I. Serrano-Pedraza (Institute of Neuroscience, Newcastle University, Newcastle upon Tyne, United Kingdom).

When the stimuli presented to each eye cannot be fused, awareness alternates between the two monocular images. This perceptual alternation is generally assumed to involve interocular suppression and adaptation, but little is known about how one image initially overcomes the other. Plausible explanations are: asymmetric interocular suppression, interocular differences in adaptation rate, or a random process. The former two imply a clear perceptual bias favoring the selection of one eye's input, whereas no differences would appear if the latter explanation is correct. Individual biases in onset rivalry were evident in 6 out of 8 observers tested, and they have also been reported previously (Carter Cavanagh, *PLoS ONE* 2, 2007). What is the explanation for these biases? We obtained the monocular contrast sensitivity for each participant, and there were no differences between the eyes. Then we assessed the contrast sensitivities when there was a mask on the other eye with contrast 30% higher than the baseline threshold. The results show that thresholds increased in a differential manner for each eye. These interocular differences in masked thresholds were negatively correlated with the proportion of times participants selected one or the other eye (i.e., lower thresholds during dichoptic masking increased the probability of selecting that eye). These findings suggest that dichoptic masking and the processes responsible for initiating rivalry are mediated by the same mechanisms. As dichoptic masking involves interocular suppression in V1 (Sengpiel & Vorobyov, *Journal of Neuroscience*, 25, 2005), our results suggest that reciprocal inhibition in the primary visual area determines the onset of binocular rivalry, and that these inhibitory connections are biased.

Financed by SEJ2007-61039/PSIC

Illusory reverse-motion caused by converging moving objects

A. Maiche (Departament de Psicologia Bàsica, Evolutiva i de l'Educació, Universitat Autònoma de Barcelona, Spain), R. Budelli (Sección Biomatemática, Facultad de Ciencias, Universidad de la República, Montevideo-Uruguay) and L. Gómez-Sena (Laboratorio de Neurociencias, Facultad de Ciencias, Universidad de la República, Montevideo-Uruguay).

In recent works (including a companion presentation to this conference), we proposed the existence of a facilitation wave that could be the basis of motion perception. A mechanism of pre-activation of neighboring areas of the cortex by a moving object seems to be a good candidate to explain our main result: The time needed to detect a target is diminished when its presentation is preceded, in a certain spatio-temporal window, by a moving object that would converge on the same spot. Based on this finding, we designed a specific stimulus configuration that induces a powerful motion illusion: We perceive apparent motion from two simultaneous targets (Gaussian patches) when one of them is preceded by a primer motion. This is probably due to a facilitation wave that is consistent with neurophysiology data concerning long-range horizontal connections, supporting the influence of these connections on motion perception. We explore this effect in greater depth by asking subjects to indicate the perceived direction of the apparent motion of two targets while the time between them was varied systematically according to the constant stimuli method. Targets are always preceded by moving Gabor patches. One of them arrives near the location where one of the targets would appear (primer), and the other moving Gabor patch stops further away from the second target. Results show that the advance in time of the perceptual appearance of the target provoked by the primer is about 40 milliseconds. This is probably due to the fact that the primer facilitates the spot where the target would appear. Thus, if the primer affects the first target, the direction of apparent motion is more clearly perceived; but if the primer affects the second target, the apparent motion is perceived in the reverse direction. This illusory reverse-motion supports the hypothesis that there is a further facilitation wave preceding any moving object and this allows us to make some predictions about the perceived duration of targets affected by motion. We expect to present herein some preliminary data about predicted overestimation of duration of targets when they are preceded by motion sequences.

Three factors in visual complexity

E. Munar and M. Nadal (Human Evolution and Cognition, associated group to IFISC (UIB-CSIC)).

Visual complexity is a very complicated concept. Our perspective in this study was from the observer's viewpoint and not as a quantifiable property of the visual array. The hypothesis refers to the relations between different features of visual complexity. We selected seven complexity dimensions we believed could relate to different aspects of visual complexity: unintelligibility of the elements, disorganization, amount of elements, variety of elements, asymmetry, variety of colors, and three-dimensional appearance. Participants were asked to rate each stimulus on a 1 to 9 Likert scale for each scale separately. Factor analysis was performed to assess the relations among the seven complexity dimensions. Factor 1 received high loadings from complexity dimensions of amount of elements, variety of elements, variety of elements, and three-dimensional appearance. Factor 2 received high loadings from dimensions unintelligibility of the elements and disorganization. Additionally, dimension asymmetry showed unsatisfactory relations with both factors and formed Factor 3. We refer to these factors as Elements, Organization, and Asymmetry. Thus, our results add further support to the idea that two or three processes contribute to the formation of subjective visual complexity. Probably, the most important process is the determination of the number and variety of elements. The second process refers to the difficulty with which the elements are identified and organized into a coherent scene. Although previous studies have subsumed asymmetry within organizational processes, our results showed that this was not an adequate solution for our data, and hence, we chose to include it as a separate factor.

SYMPOSIUM 3: VISUAL PERCEPTION

Moderator: Miguel Castelo-Branco

Ambivalence and ambiguity in facial expression of emotions

M.A. Cahill and M.A. Aguilar (Universidad Complutense de Madrid).

Perception of mixed emotions in pure and mixed facial expressions was studied. In Experiment 1, participants gave independent evaluations of positive and negative valence and of familiarity of a set of pure (anger, disgust, fear, happiness, and sadness) and artificially constructed mixed expressions. Mixed expressions were composed of a happy upper or lower half face and an upper or lower half face corresponding to each of the four negative emotions. Mixed expressions were evaluated as being more ambivalent (positive and negative simultaneously) than pure expressions and this was specially so for expressive mixes containing a happy mouth. In Experiment 2, the extent to which each expression was recognized as showing different emotions at the same time (emotional ambiguity) was studied. Participants had to identify the expression of each face, choosing from various terms that referred to basic emotions and cognitive states (surprise, strangeness). Participants were significantly more accurate in identifying

pure than mixed expressions. Moreover, they identified two different emotions more frequently in mixed than in pure expressions, although, especially in the case of fearful faces, two emotions were also identified in some pure expressions. In most cases of double identification, “surprise” appeared as one of the expressions identified. In the case of mixed expressions, this occurred with faces showing happiness in the lower half face. In Experiment 3, sensitivity of expression identification to the immediately preceding linguistic context was studied. Each expression was preceded in different trials by the name of different emotions that acted as a prime, and participants had to indicate to what extent the face showed that specific emotion. Even though the resulting emotion profiles showed in general that mixed expressions were more ambiguous than pure expressions, the participants also perceived different emotions in the pure expressions, especially in the case of fearful expressions, which were recognized as showing fear and surprise at the same rate. Our results show a considerable degree of ambivalence and ambiguity in the perception of emotional expression and suggest that the immediate context in which an expression takes place can have a profound influence on the affective meaning assigned to that expression.

Supported by: Ministerio de Educacion y Ciencia de España (ref: SEJ2006-01576/PSIC).

Modulation of perceptual processing by the affective meaning of faces: An ERP (event-related potentials) study

B. Valdés-Conroy, L. Aguado, T.D. Risco, S. Rodríguez and F.J. Román (Universidad Complutense de Madrid, Spain).

Using an associative learning task, we studied how facial expression and acquired emotional value modulate the face-specific N170, an event-related potential (ERP) characteristic of visual perception of faces. Participants had to learn and associate the identity and expression (angry, happy, or neutral) in a series of neutral-expressive pairs of faces. Our results show modulation of the N170 over occipito-temporal sites due to facial expression (happy and angry vs. neutral). Furthermore, this modulation was also sensitive to expressive value (happy vs. angry). Modulation of the N170 was also observed by the identity of faces, depending on the associated expression. An attenuation of the N170 response to angry faces after repetition of exposures was also observed. Similar results were obtained at fronto-central locations within the N170 time window both for expressive and neutral associated faces. These results show the effect both of intrinsic and acquired affective meaning of faces on early perceptual face processing and have important implications for current theories on face processing and the relationship between emotion and cognition.

This study was supported by the project SEJ2006-01576/PSIC.

Neural correlates of ambiguous face detection

J. Castelhana (Visual Neuroscience Lab, IBILI, University of Coimbra, Portugal), J. Rebola (Visual Neuroscience Lab, IBILI, University of Coimbra, Portugal), E. Rodríguez (Pontificia Universidad Católica de Chile, Escuela de psicología, Santiago, Chile and Max-Planck for Brain Research, Frankfurt am Main, Germany), B. Grawe (Visual Neuroscience Lab, IBILI, University of Coimbra, Portugal) and M. Castelo-Branco (Visual Neuroscience Lab, IBILI, University of Coimbra, Portugal).

The visual processing of faces and objects has received considerable attention in recent years. We have used event-related technology (EEG/ERP) to help identify the processes underlying the emergence of a coherent face percept under ambiguous conditions. It has been hypothesized that these processes relate to neural synchrony in the gamma band (30-80 Hz) close to the moment of perception. Here, we aimed to investigate this hypothesis by dissociating stimulus from perceptual factors in the face perception process. EEG data from 21 subjects were recorded (64/128 channel Quick-Cap, NeuroScan). During the task (Experiment 1), subjects viewed 12 s movies of rotating ambiguous Mooney faces and were instructed to report the moment of face detection by pressing a button. The data were segmented into epochs locked to the response (moment of perception). Decision locked event-related potentials (ERP) were obtained, and time-frequency as well as phase synchrony analysis on the gamma band range were carried out across distinct electrode sets. In Experiment 2, face stimuli comprised one frame. Two additional noise frames were added for backward/ forward masking. The behavioral results of Experiment 1 show a mean response time of 4.23 ± 2.98 s which corresponds approximately to an 85° angle of stimulus rotation—very close to the horizontal meridian, suggesting holistic processing of the stimuli (mean accuracy 90.78 ± 9.36 %). In Experiment 2, jittered face stimuli were always perceived as more salient in the absence of forward masking. Spatially distributed activation in the high gamma band (60-70 Hz) was observed in the time-frequency plots, and synchrony bursts (30-45Hz) were also observed, close to the moment of detection. Similar findings were obtained in Experiment 2. In conclusion, our data support that gamma-band activity is related to perceptual decision and not only to perception of low level features *per se*.

Funding acknowledgments: FCT Proj. PTDC_PSI_67381_2006.

Perception of radial motion relies on detecting spatial displacements

C. de la Malla and J. López-Moliner (Vision and Control of Action (VISCA) lab.; Institut de Recerca en Cervell, Cognició i Conducta; Universitat de Barcelona).

It has been shown that unconfounding the use of motion from spatial displacement information is possible for lateral motion (Nakayama & Tyler, 1981). Here, we determine whether this is also possible for radial motion. For this purpose, we adapted the paradigm previously used by Nakayama and Tyler to obtain detection thresholds for lateral and radial motion by adjusting the displacement amplitude of random dot patterns. In agreement with previous studies, we found that the detection of lateral motion relies on either motion or displacement, depending on the range of temporal frequency and stimulated area. However, the detection of radial motion was always consistent with detecting a displacement threshold.

Learning to listen through another person's ears: A study of learning curves using non-individualized Head-Related Transfer Functions

J.A. Santos (Departamento de Psicologia, Universidade do Minho, Portugal), C. Mendonça (Departamento de Psicologia, Universidade do Minho, Portugal), C. Murteira (Departamento de Psicologia, Universidade do Minho, Portugal), J.P. Ferreira (Centro de Território Ambiente e Construção, Universidade do Minho, Portugal), P. Dias (Departamento de Electrónica, Telecomunicações e Informática / IEETA, Universidade de Aveiro, Portugal) and G. Campos (Departamento de Electrónica, Telecomunicações e Informática / IEETA, Universidade de Aveiro, Portugal).

Virtual Reality (VR) environments traditionally aim at visual immersion, but the development of convincing VR simulations requires that other senses also be considered. Aural immersion, or auralization, is the natural next step in the evolution of VR systems. Auralization relies on anechoic sound source recordings processed to mimic the influence of the acoustic propagation environment (absorption, reflections, reverberation). The listener's head, pinna, and torso have a crucial effect that can be described by Head-Related Transfer Functions (HRTF), different for each individual. Earphone presentation makes it possible to auralize sounds processed using HRTF other than the listener's own. In our study, we used non-individualized HRTF, obtained from mannequins equipped with ear microphones. Subjects were asked to indicate the perceived source position for sounds processed to simulate different azimuth angles: from 0° (front) to 90° (right). Several sessions were carried out for each participant. The goal was to find how their localization accuracy varied over time, as it was hypothesized that exposure would induce a learning effect. Preliminary data indicate that 3D localization does indeed improve with exposure but the learning curves evidence a plateau effect, beyond which further exposure does not lead to additional improvement.

Supported by HFSP, DFG, EC FP6 project COBOL and FP7 project SEARISE, and the Hermann und Lilly Schilling Foundati.

SYMPOSIUM 4: COLOUR VISION

Moderator: Julio Lillo

Color categories in infancy and early childhood

A. Franklin (Department of Psychology, University of Surrey).

The origin and nature of color categories in language and cognition has been the concern of researchers from a range of disciplines such as psychology, anthropology, cognitive science, linguistics and philosophy for many decades. One major issue is whether the color spectrum is arbitrarily carved up into categories, or whether there are universal constraints on where these categories form. In support of the argument that there are constraints on how language categorizes color, there is converging evidence for categorical responding to color before the acquisition of color terms (e.g., Franklin & Davies, 2004; Franklin, Pilling & Davies, 2005). This talk will outline a series of studies that have attempted to establish the nature and underlying mechanisms of these prelinguistic categories. In one study, the event-related-potential technique is used to identify the neural markers and underlying mechanisms of categorical responding to color in 7-month old infants (Clifford, Franklin, Davies & Holmes, 2009). Another set of studies have considered how pre-linguistic color categories are lateralized in the human brain. Whilst categorical responding to color in adults is stronger in the left hemisphere (e.g., Gilbert, Regier, Kay & Ivry, 2005), in 4-6 month-old infants, the category effect is lateralized to the right hemisphere (Franklin, Drivonikou, Bevis, Davies, Kay & Regier, 2008). The change in lateralization appears to be related to language as there is a right hemisphere bias for toddlers who do not know the terms for the relevant categories, yet a left hemisphere bias for those who have learnt the terms

(Franklin, Drivonikou, Clifford, Kay, Regier & Davies, 2008). These findings suggest that there is a right-hemisphere substrate for categorical responding to color that is replaced or suppressed by the mechanisms of language once color terms are learnt. The implications for our understanding of the origin and nature of color categories in language and cognition are discussed.

A categorical color effect in the brown-orange boundary

J. Lillo (Dep. Psicología Diferencial y del Trabajo. Universidad Complutense de Madrid. Spain), A. Al-Rasheed (Dep. Psychology. University of Surrey. Guilford. United Kingdom) and H. Moreira (Dep. Psicología. Universidad Cardenal Cisneros. Madrid. Spain).

A categorical color effect appears when, for similar inter-stimuli distance, a larger perceptual change is detected for stimuli belonging to different categories (inter-category change) than for stimuli belonging to the same category (intra-category change). Our research framework was the brown-orange transition. All the stimuli we used had the same chromatic coordinates ($u' = 0.32$; $v' = 0.53$), but they varied in luminance. In the first experiment, stimuli were named either brown or orange (forced choice) and were presented on two different backgrounds (articulated or non-articulated). Background type influenced the luminance value where the brown-orange transition appeared. Its value was larger for the articulated background. This result agrees with the predictions derived from Gilchrist's anchoring theory on lightness perception and indicates that L^* calculus must consider the background type on which a stimulus is presented. A second experiment detected a categorical perception effect using a visual searching task. This task required observers to locate (right or left) a target stimulus. Shorter times and fewer errors appeared when target and distracters belonged to different categories (inter-category change). However, and in contrast to other works, no evidence of laterality was found (the categorical effect appeared both for stimuli on the right and on the left).

Supported by PSI2008-04166.

The relationship between categorical color perception and shift towards prototype

I. Davies (Department of Psychology, University of Surrey, UK) and E. Özgen (Department of Psychology, University of Bilkent, Ankara, Turkey).

Discrimination among stimuli is affected by systematic biases in detecting the precise properties of stimuli. These biases include: central tendency (Hollingworth, 1910); shift towards prototype (STP; Huttenlocher et al., 2000); and category truncation (Huttenlocher, 2007). These biases could underlie perceptual warping (Harnad, 1987) evidenced in categorical perception (CP): better discrimination of stimuli from different categories than of stimuli from the same category. However, Pilling et al., (2003), using a same-different discrimination task, found clear evidence of both central tendency and categorical perception, but argued that the two were independent. Here, based on reanalysis of Özgen and Davies's learning experiments and a matching-to-sample memory task, we report evidence that STP could be the basis of newly learned CP. In Experiment 1, subjects practiced a successive same-different task on stimuli differing in hue or lightness, where the average of the stimuli and the category prototype(s) coincided. There was clear evidence of bias towards the coincident point. In subsequent experiments, subjects learned to divide either blue or green into two new categories, with the new boundary at the category prototype. After training, in discrimination tests using a successive same-different task, the average of the test stimuli again coincided with the blue or green prototype, and untrained subjects showed the same bias towards the center as in Experiment 1. Trained subjects showed enhanced discrimination across the new boundary (CP), but also showed bias in the opposite direction from the test-stimuli average. This change in direction of bias could be due to bias towards newly abstracted prototypes. In the matching-to-sample task, stimuli ranged from yellow to purple with the average at the blue-green boundary. Judgments showed both a bias towards prototypes, and a bias towards the center of the test range. The relationship among the various perceptual biases and CP is discussed.

Color categorical perception and color metrics

Oliver Wright (Bilkent University, Ankara, Turkey).

Color categorical perception (CP) refers to the idea that pairs of colors belonging to different categories can be discriminated faster and/or more accurately than pairs of colors belonging to the same category. Unlike studies investigating the CP of speech sounds, which are usually specified using a physical metric, research into color CP typically uses stimuli specified in perceptual metrics. The most widely used perceptual color metric and the one considered to show the highest degree of perceptual uniformity is the Munsell system (Munsell, 1905; Newhall, Nickerson, & Judd, 1943). It is suggested here that specifying stimuli using perceptual color metrics presents some significant challenges for research into color CP. First, much experimental evidence for color CP rests on the

assumption that the perceptual metrics used are uniform. But research suggests that this is not the case (e.g., Indow, 1988; Indow & Kanazawa, 1960). This implies that much of the evidence for color CP can alternatively be interpreted as evidence of irregularities in the color metrics used. Second, there seems to be no reason to suppose that color CP is not “built into” the color metrics used. This presents additional difficulties for some theories of color CP.

Category training affects color discrimination

G.V. Drivonikou (University of Surrey, U.K.), A. Clifford (University of Surrey, U.K.), A. Franklin (University of Surrey, U.K.), E. Özgen (Bilkent University, Ankara, Turkey) and. I. Davies (University of Surrey, U.K.).

Categorical perception (CP) of color is shown when discrimination is faster and more accurate for two colors from different color categories than for two colors from same categories, even when chromatic separation sizes are equated. There is indirect evidence that color CP may be learned. For instance, CP is only shown if the category boundary is marked in the subject’s language (e.g., Roberson et al., 2000). There is also evidence that CP can be induced across a newly learned boundary (Özgen & Davies, 2002), as indicated by performance in a subsequent same-different discrimination task. However, as the latter measure could be influenced by high level processes such as labeling or other forms of on-line language influence, here, we replicate and extend Özgen and Davies’ category learning study in an attempt to pinpoint the nature of the changes underlying category learning. Subjects learned to divide the green region into two new categories “yellow-green” / “blue-green” over four days. Together with a control group, they were then tested on green and blue regions, using a target detection task where colored targets were presented on same- or different-category backgrounds. The trained group showed CP across the new boundary on the target detection task, whereas the controls did not. Moreover, CP was stronger for right visual field targets than for left visual field targets (cf. Drivonikou et al. 2007). The results suggest that learning to categorize stimuli alters their perception at an early stage of visual processing. The findings are related to the debate about the nature and origin of color CP.

Chromatic diversity of indoor scenes rendered with CIE illuminants and white LEDs

J.M.M. Linhares, P.E.R. Felgueiras, P.D. Pinto and S.M.C. Nascimento (University of Minho, Department of Physics, Braga, Portugal).

The color quality of a light source is typically evaluated by the color rendering index (CRI), a quantity with well-known limitations. The goal of this work was to evaluate the quality of lighting by estimating instead the chromatic diversity light sources produce in indoor scenarios, using CIELAB and DIN99 color spaces. Reflectance spectra of objects typically found in indoor scenes (such as books, colored fabrics, children’s toys, fruits, indoor plants, among others) were obtained using an hyperspectral imaging system with a low noise cooled digital camera with a spatial resolution of 1024 (H) x 1344 (V) pixels and a fast tunable filter with an infrared filter in front of the lens. Images were acquired from 400 to 720 nm in 10 nm steps. Care was taken to avoid shadows and multiple reflections. Chromatic diversity was estimated for 60 illuminants, 55 CIE illuminants, and 5 LED light sources (Luxeon, Philips Lumileds Lighting Company, USA) by computing the CIELAB and DIN99 color volumes of the objects and by counting the number of non-empty unitary cubes of the segmented volumes. A large variation in chromatic diversity was found across illuminants; in addition, a low degree of correlation between the number of discernible colors and the CRI of the illuminant was also found. The best illuminant was CIE FL3.8, producing about 7.5% more colors than CIE illuminant A and it was 8% better than D65 if CIELAB was considered, and CIE FL11 produced about 7.4% more colors than CIE illuminant A and was 4% better than D65 if DIN99d was considered. These results suggest that normal observers may benefit from a careful choice of the illuminant to maximize the chromatic diversity perceived in indoor scenes and this choice is not necessarily based on the CRI.

This work was supported by the Centro de Física of Minho University, Braga, Portugal, by the Fundação para a Ciência e a Tecnologia (grants POSC/EEASRI/ 57554/2004 and POCTI/EAT/55416/2004).

João M.M. Linhares was supported by grant SFRH / BD / 35874 / 2007.

SYMPOSIUM 5: TEMPORAL ASPECTS OF PERCEPTION

Moderator: Joan López-Moliner

The role of temporal processing and feedback in visual attention and awareness TAP1

S.L. Macknik (*Barrow Neurological Institute of St. Joseph’s Hospital and Medical Center, USA.*)

The mammalian visual system includes numerous brain areas that are profusely interconnected. With few exceptions, these connections are reciprocal. Anatomical feedback connections in general outnumber feed-forward connections, leading to widespread speculation that feedback connections play a critical role in visual awareness. However, temporal processing evidence from physiological experiments suggests that feedback plays a modulatory role, rather than a driving role. I will discuss theoretical constraints on the significance of the anatomical numerical advantage of feedback, and describe theoretical limits of its potential physiological impact. These restrictions confine the potential role of feedback in visual awareness and rule out some extant models of visual awareness that require a fundamental role of feedback. I will propose that the central role of feedback is to maintain visuospatial attention, rather than visual awareness. These conclusions highlight the critical need for experiments and models of visual awareness that control for the effects of attention.

Reaction time used as a mean to assess perceptual latencies in cortical motion processing

L. Gomez-Sena (*Laboratorio de Neurociencias, Facultad de Ciencias, Universidad de la República, Uruguay*), R. Budelli (*Laboratorio de Neurociencias, Facultad de Ciencias, Universidad de la República, Uruguay*) and A. Maiche (*Laboratori de Percepció i Psicofísica, Universitat Autònoma de Barcelona, Espanya*).

In a previous work, we showed that a moving object (primer) whose trajectory intersects the trajectory of another object (target) increases the magnitude of the flash lag illusion. Based on those results, we argued that a pre-activation mechanism could be involved in the generation of the illusion, provided by the combination of the feed-forward sensory input and intra-cortical horizontal connections, which result in a reduced perceptual latency. A consequence of this mechanism would be that the reaction time (RT) to the detection of a target should be less when it is preceded, in a certain time window and spatial surround, by a moving object that converges to the same spot. By inducing motion illusion with short sequences of Gabor patches and varying the spatial and temporal relations between them and the target, we showed that the RT is, indeed, modulated in a manner consistent with the pre-activation mechanism. In the limit, when just one primer is present, we obtain the configuration studied by Polat and Sagi (1993) where two Gabor patches were presented flanking a central target. They showed that the flankers, depending on the lateral distance and timing, reduce the threshold for target detection. We carried out similar experiments using RT and we obtained similar qualitative and quantitative results, thus gaining further support to our approach. Interestingly, Polat and Sagi relate their findings to contour integration and we relate ours to motion perception. Naturally, the same circuitry is activated in both cases, and the outcome of the processing will depend on contextual elements and the reciprocal connections with other areas. A neural pattern will prevail, emerging from those dynamic interactions, which will ultimately determine the effective percept.

Physiological influences on the rate of subjective time: The case of hypercapnia

D.T. Field (School of Psychology Clinical Linguistic Sciences, University of Reading, UK and Centre for Integrative Neuroscience Neurodynamics, University of Reading, UK), C. Whyborn (School of Psychology Clinical Linguistic Sciences, University of Reading, UK) and C.A. Scott (School of Psychology Clinical Linguistic Sciences, University of Reading, UK).

For short intervals, subjective time is a linear function of real time, and this is often explained by proposing the existence of a pacemaker in the brain that generates timing pulses. As a biological mechanism, the rate at which the pacemaker produces pulses is thought to be influenced by various physiological factors, such as arousal, the actions of various drugs, and body temperature. For example, a reduction of body temperature slows the rate of subjective time, which is thought to be due to a reduced rate of pulse production from the pacemaker. Likewise, an increase in body temperature increases the rate of subjective time due to an increase in the rate of pulse production. We present the first investigation of the effects of a hypercapnia on the rate of subjective time. Hypercapnia is a physiological change induced either by breathing air with a higher than normal percentage of CO₂ or by breath holding (used here). Hypercapnia changes the physiological environment of the brain by increasing the proportion of CO₂ in the blood, and by producing vasodilation leading to an increase in cerebral blood volume. In the experiment, participants made verbal estimates of auditory stimuli with durations in the range 150 msec to 950 msec. Auditory stimuli were either continuous filled tones, or two brief clicks defining an empty interval. Previous research has indicated that filled intervals appear longer than unfilled intervals (the so called "filled duration illusion"), and this has been explained by suggesting that the pacemaker runs faster for filled than for unfilled intervals (Wearden et al., 2007). We replicated the filled duration illusion under conditions of normocapnia, and found that hypercapnia abolished the filled duration illusion by shortening estimates of filled intervals relative to normocapnia while leaving estimates of unfilled intervals unchanged relative to normocapnia. A plausible interpretation of this pattern of results is that hypercapnia prevents the pacemaker from running at the higher rate it normally runs at for filled durations.

Time in motion: Speed channels and the color-motion asynchrony

J. López-Moliner (Vision Control of Action, Institut de Recerca en Cervell, Cognició i Conducta, Universitat de Barcelona) and D. Linares (School of Psychology, University of Sydney, Australia).

Motion has intricate effects on time-related judgments. One of the most well-known effects is the color motion asynchrony illusion (CMA): When color and motion direction changes occur in synchrony at a high rate, we perceive these changes as occurring asynchronously. We have suggested (Linares et al., 2006 *Journal of Vision*, 6, 974-981) that the opponency mechanisms that underlie motion perception are the cause of the CMA. Assuming little cross-talk between the processing of slow and fast speeds (van der Smagt et al., *Nature Neuroscience*, 2, 585-596), our interpretation would gain support if the perceived CMA was reduced by using different speeds (e.g., slow-downwards and fast-upwards). To test this, observers had to pair color (red/green) with direction (upwards/downwards) of moving plaids. We sampled ten phase differences between color and motion within a cycle of 600 ms in three speed conditions: slow-slow, fast-fast, and slow-fast. The results confirmed a perceptual effect that can be acknowledged by eye inspection only: the reduction of the CMA (about 40 ms) when the speeds in the two directions were different (slow-fast). These results not only provide support for the role of motion opponency in the CMA illusion, but also add further evidence for slow and fast speed segregation.

Supported by SEJ2007-60488/PSIC from the Spanish Government.

SYMPOSIUM 6: TOUCH: BEHAVIORAL DATA AND BRAIN ACTIVATIONS

Moderator: Soledad Ballesteros y José Manuel Reales

A descriptive model of horizontal gradient and oblique effect in the haptic parallelity task

M. Fernández and D. Travieso (Dpto. Psicología Básica, Facultad de Psicología, Universidad Autónoma de Madrid).

Previous research on the properties of haptic space has shown systematic deviations when performing parallel matching tasks, depending on horizontal distance and orientation of the bars, which are described as deformations of Euclidean space, or better, as non-Euclidean space. In addition, a haptic oblique effect, that is, increments in the deviations when matching oblique orientations, has been shown. There are several models developed to describe this striking effect. Cuijpers, Kappers, and Koenderink (2003), and Kappers and Volvic (Kappers & Volvic, 2007; Volvic & Kappers, 2008) understand haptic spatial performance as a representational problem: in these matching tasks, the perceived orientation of a reference bar is represented spatially in some frame-of-reference, and then has to be reproduced in the orientation of a test bar. Experimental results in parallel matching tasks show that performance can neither be described with an egocentric frame-of-reference nor an allocentric one, but with a complex composition of both reference frames. There are two main weaknesses in the above-mentioned models. First, the general descriptive model by Cuijpers and cols. (o.c.) does not consider the strong individual differences in performance, whereas Kappers and Volvic (o.c.) present an additive model with a strong inferential level based on representations. In addition, none of these models give account of the oblique effect that is systematically found in the task. Our study presents data of the standard haptic parallelity task for 5 subjects and develops an additive linear model of performance, including the oblique effect. Through the maximum likelihood method, subjects are assigned to 1 out of 3 hierarchical models, which reflects the amount of the deviations as well as the strength of the oblique effect in performance.

This research was supported by grant HUM2006-11603-C02-02 of the Spanish Ministry of Science and Innovation.

The haptic horizontal-vertical illusion across adulthood

M.A. Heller (Northwestern Illinois University, USA), J. Mayas (Universidad Nacional de Educación a Distancia, Madrid, Spain), J.M. Reales (Universidad Nacional de Educación a Distancia, Madrid, Spain) and S. Ballesteros (Universidad Nacional de Educación a Distancia, Madrid, Spain).

The haptic vertical-horizontal illusion is not dependent on visual status or visual imagery, as similar illusory overestimation of the vertical occurs independently of the visual status (early blind, later blind, and low vision) of the perceiver (Heller et al., 2003). The present study was conducted to investigate whether this spatial misperception remains stable or increases across adulthood and old age. In the present study participated young adults (22-40 years), middle aged adults (41-59 years), and older adults (60-75 years). Participants examined unimanually or bimanually in a counterbalanced order 5 horizontal and 5 vertical curved shapes varying in size, and estimated size (horizontal/vertical) using two slide rules. The results suggest that the illusion is independent of age: (a) all groups

overestimated the vertical; (b) the illusion did not change as a function of stimulus size; and (c) the mode of exploration (unimanual or bimanual) did not affect the illusion.

Neural correlates of textured stimuli using a discrimination task: An electrophysiological study

F. Muñoz, M. Sebastián, J.M. Reales and S. Ballesteros (Universidad Nacional de Educación a Distancia, Madrid, Spain).

In the present study, we investigate the cortical modulations involved in tactile roughness perception using a tactile texture discrimination task. Twenty-two volunteers were presented with rough and smooth surfaces using a classical odd-ball paradigm (target, $P = 0.2$ and standard, $P = 0.8$). The textured surfaces were mounted on a specifically designed apparatus (the Haptic Spinning Wheel), which moved at a constant velocity under the static index fingertip of the perceiver. Participants mentally counted the number of targets perceived. The ERP results showed an increased P300-like component for attended compared to unattended stimuli. A time-frequency analysis yielded strong synchronization in the theta band (3-7 Hz) at an early time window (around 200 ms), followed by desynchronization in the alpha band (8-12 Hz) along the 400-800 ms interval. LORETA source analysis showed higher neural activation when processing targets compared to standard stimuli at somatosensory, occipital, and frontal cortices. These findings suggest that top-down processes may modulate the information processing of tactile stimuli by increasing and decreasing particular brain oscillations in different cortical areas.

This research was supported This work was supported by a grant from the European Community: NEST-2005-Path-IMP grant 043432 (SOMAPS) to S.B.

SYMPOSIUM 7: HEARING AND SPEECH

Moderator: Luís Enrique López-Bascuas

The role of acoustic scale in the perception of musical notes and instruments

R. Paterson (Centre for the Neural Basis of Hearing, Department of Physiology, Development and Neuroscience, University of Cambridge, UK).

This talk is about the sounds made by musical instruments and how we perceive those sounds. It is intended to explain the basics of musical note perception; that is, why instruments come in families; what determines “register” within families; and why we hear distinctive differences between members of a given instrument family—even when they are playing the same note. On the surface, the answers to these questions may seem obvious; one could say that brass instruments all make the same kind of sound because they are all made of brass, and the different members of the family sound different because they are different sizes. But there is a deeper explanation involving three acoustic properties of musical sounds as they occur in air. The talk describes these properties (with audio demos) and explains why they are particularly useful in (a) summarizing the physics of note production by instruments, on the one hand, and (b) explaining the dimensions of musical note perception, on the other hand.

Auditory discontinuities and the perception of vowel height in Spanish back vowels

L.E. López-Bascuas (Universidad Complutense de Madrid, Spain).

A wide-spread approach to the study of speech perception entails binding phonological categorization to some sort of auditory discontinuities. Consistent with this view, some authors made an attempt to connect the “center of gravity effect” (CGE) (Chistovich & Lublinskaya, 1979) to the perception of vowel height, suggesting that the [+/- high] boundary might be based around a “quantal” region of the auditory space. In particular, Syrdal and Gopal (1986) found that productions of American vowels could be divided into the phonological categories [+/- high] by a boundary at 3-3.5 Bark F1-f0 distance (which corresponds to the limit of integration of the CGE). Later, in a perceptual study using front vowels, Hoemeke and Diehl (1994) found that the boundary for the phonological feature [+/- high] was indeed at 3-3.5 Bark F1-f0 distance. However, Fahey and López-Bascuas (1994) and López-Bascuas et al. (1995) found that the 3-3.5 Bark F1-f0 boundary does not seem to be effective for Spanish front vowels either in production or in the perception of the [+/- high] phonological contrast. Moreover, in English, results obtained with back vowels do not seem to support clearly the 3-3.5 Bark hypothesis (Fahey, Diehl, & Traunmüller, 1996). In this study, our previous results with front vowels are extended to Spanish back vowels. F1 and f0 were measured from Spanish vowel productions in order to estimate the F1-f0 Bark value that separates the [+/- high] back vowel categories. Our acoustic analyses agree well with our perceptual results and do not support the

hypothesis of a 3-3.5 Bark F1-f0 integrator playing a significant role in the psychological organization of the [+/-high] distinction.

Supported by SEJ2006-11955.

Psychoacoustical verification of current theories on the generation of human distortion product otoacoustic emissions

E.A. Lopez-Poveda and P.T. Johannesen (Instituto Neurociencias de Castilla y León, Universidad de Salamanca, Salamanca, Spain).

When two pure tones (or primaries) of slightly different frequencies (f_1 and f_2) are presented to the ear, new frequency components are generated by the nonlinear interaction of the primaries in the cochlea. These new components can be heard and emitted from the cochlea back to the ear canal. The level of the $2f_1 - f_2$ emitted distortion is regarded as an indicator of the physiological state of the cochlea. This indicator, known as the distortion product otoacoustic emission or DPOAE, is most sensitive when the primaries have levels that evoke maximal-level DPOAEs. Efforts have been directed to obtain these levels empirically but the optimal primary-level combination is still controversial. The controversy could be clarified by elucidating the cochlear mechanical conditions that maximize DPOAE levels. The current view is that maximal-level DPOAEs occur for primaries producing equal excitation at the cochlear place tuned to f_2 , but this conjecture cannot be tested directly in living humans because it is impossible to record cochlear motion while monitoring ear canal DPOAE levels. Here, the conjecture is tested using psychoacoustical methods like those used to infer human peripheral compression. Results support the current view for f_2 frequencies of 1 and 4 and for levels below around 65 dB SPL. The present approach may be extended to test other uncertainties on the generation mechanisms of human DPOAEs.

Supported by MEC (BFU2006-07536) and The Oticon Foundation.

Vowel formant structure and speaker identification. A perceptual study

M.J. Albalá (Consejo Superior de Investigaciones Científicas), E. Battaner (Universidad Rey Juan Carlos), J. Gil (Consejo Superior de Investigaciones Científicas), J. Llisterri (Universidad Autónoma de Barcelona), M. Machuca (Universidad Autónoma de Barcelona), V. Marrero (Universidad Nacional de Educación a Distancia), C. de la Mota (Universidad Autónoma de Barcelona), M. Marquina (Universidad Autónoma de Barcelona) and A. Ríos (Universidad Autónoma de Barcelona).

This study belongs to VILE Project (Acoustic and Perceptual Study of Variation within and between Speakers in Spanish), whose goal is to validate—acoustically and perceptually—relevant cues to establish the individuality of speakers, in order to apply the results to automatic speech recognition and to judicial practice (Battaner et al., 2003; Marrero et al., 2003). On this occasion, we shall present the results of a natural vowel 2IAX discrimination test. Stimuli were extracted from the acoustic database used in VILE-1st part. 40 pairs (4 vowels —/i,e,a,o/—, 2 accents —stressed/unstressed—, and 5 acoustic parameters—F0/F1/F2/F3/F4), as similar as possible in all acoustic parameters except for the one under study were presented to 46 judges (34 women and 12 men) aged 18-35. Stimuli were delivered via headphones in a soundproof radio cabin using Paradigm software (Lopez-Bascuas et al., 1999). Results show 94.53% of hits when stimuli were the same, and 72.97% for different stimuli. Concerning the four vowels, the proportion of errors was 28.47% for /o/, 27.01% for /a/, 26.48% for /e/, and 26.14% for /i/. As for the acoustic parameters, the largest error rate corresponds to F4 (39.97%), F3 (35.35%), and F1 (31.09). On the contrary, F0 ratio is only 13.73% and F2 14.70%. Thus, our findings seem to indicate the prevalence of F0 and F2 as main cues for speaker identification across different vowel categories. In the conclusion, implications of these results for future research are discussed.

Supported by the Ministerio de Ciencia y Tecnología (BFF2001-2551) and the Ministerio de Educación y Ciencia (HUM2005-06980).

Asymmetry in auditory priming: Evidence from the perception of words, sounds, and talkers

J.G. Alvarez (Universitat Jaume I. Castellón de la Plana, Spain) and C.T. McLennan (Cleveland State University, OH, USA).

Marsolek and colleagues have found support for two dissociable and parallel neural subsystems underlying the recognition of written words, visual objects, and visual shapes: an abstract-category subsystem that operates more effectively in the left cerebral hemisphere (LH), and a specific-exemplar subsystem that operates more effectively in

the right cerebral hemisphere (RH). Thus, in the visual domain, evidence of this asymmetry exists for linguistic (words, pseudoword forms) and non-linguistic (objects) stimuli. Our work in this area was originally motivated by the idea that asymmetry may not be limited to the visual domain. Indeed, we have since obtained hemispheric differences in priming effects in the auditory domain for linguistic (spoken words) and non-linguistic (environmental sounds) stimuli. More precisely, we observed specificity effects when listeners heard auditory stimuli in their left ear (RH), but not when they heard these same stimuli in their right ear (LH). Furthermore, our more recent data on talker identification once again reveal this same pattern of asymmetry. Taken together, this consistent pattern of data from both domains may be indicative of a more general property of the human perceptual processing system.

POSTER SESSION 1

Psychophysical correlates of bias in decision-making

J.L. Pardo-Vázquez (Universidad de Santiago de Compostela, Spain), I. Padrón (Universidad de Santiago de Compostela), J. Fernández-Rey (Universidad de Santiago de Compostela), V. Leborán (Universidad de Santiago de Compostela) and C. Acuña (Universidad de Santiago de Compostela E-15705).

The sensory information required to make a decision can be inferred before obtaining evidence about it. Formally, this is the prior probability that one of the hypotheses to make a decision is true before obtaining evidence about it. How to reveal this bias and what and where are their neural correlates? To address these questions, we designed a task in which the prior probability that a given response is correct depended on a visual cue. In a discrimination reaction-time task in which they had to decide whether the length of one line (S2) was longer or shorter than another one, subjects showed a few milliseconds before (S1). Subjects communicated their decision by clicking one of two buttons of a computer mouse. A bias in the response was introduced by manipulating the prior probability that one of the responses was correct. The color of two circles displayed at both sides of the screen changed from block to block (64 trials/block): when the color was blue (low bias), S2 was shorter in 75% of the trials; when it was red (large bias), S2 was shorter in 25% of the trials; when it was gray (neutral trials), S2 was shorter in 50% of the trials. Five subjects (1 male and 4 females, ages 21-24) were tested in the task, during nine sessions each subject. Psychophysical and signal detection theory analyses suggest that this task induces and reveals the subject's bias during decision-making. Biases related to probability and decision-making significantly affect our everyday behavior. Furthermore, our task allowed us to measure cognitive responses that depend on memory traces left by visual stimuli and to study the neural correlates of these processes.

Supported by MCINN to CA and by INCITE-Xunta de Galicia (Spain) to JF-R and CA.

Does illusory surface inclination affect eye movements?

L.P Zapata (Vision and Control Action (VISCA-Lab), Facultat de Psicologia, Universitat de Barcelona), A. Aznar-Casanova (Vision and Control Action (VISCA-Lab), Facultat de Psicologia, Universitat de Barcelona) and H. Super (Vision and Control Action (VISCA-Lab), Facultat de Psicologia, Universitat de Barcelona and Institució Catalana de Recerca i Estudis Avançats (ICREA), Barcelona).

Binocular vision requires that we foveate both eyes on the relevant visual information to fuse the foveal images into one single image. Other points in space do not fall on corresponding retinal locations, and the visual system uses this binocular disparity to extract depth information. Besides binocular disparity, the visual system employs monocular cues such as texture gradients and perspective to extract depth information. In perceptual illusions, the monocular and binocular depth cues are different, which may cause a conflict for the visual system. Binocular depth cues make the eyes converge on the physical plane (screen), whereas monocular cues produce convergence on the induced depth plane. In this work, we registered eye movements in observers that looked at a point presented on different inclinations and slants of surfaces, induced by means of different gradient textures. The preliminary results suggest that there are differences in eye positions during the first 80 ms. After this initial fixation, an adjustment occurs where the eyes return to fixate at the screen plane. Thus, it appears that the oculomotor system first programs eye movement based on the induced depth perception, and only after this can the visual system correct this "depth illusion."

How does illusory surface inclination affect the accuracy of orientation judgments?

L.P Zapata (Vision and Control Action (VISCA-Lab), Facultat de Psicologia, Universitat de Barcelona), M. Moreno (Vision and Control Action (VISCA-Lab), Facultat de Psicologia, Universitat de Barcelona), H. Super (Vision and Control Action (VISCA-Lab), Facultat de Psicologia, Universitat de Barcelona and Institució Catalana de Recerca i Estudis Avançats (ICREA), Barcelona) and A. Aznar-Casanova (Vision and Control Action (VISCA-Lab), Facultat de Psicologia, Universitat de Barcelona).

A possible description of our surroundings is a spatial disposition of surfaces that differs in orientation, size, reflectance, texture, and spatial location. When the slant of the surface is known, our visual system can greatly improve the precision and accuracy of the spatial location, orientation, and distances of stimuli. To correctly perceive our surroundings, the visual system has to perform diverse geometric transformations of surfaces. Here, two of these transformations were studied. In the first task, a segment (or the end-points of a segment) is shown on an induced inclined surface (created by a texture gradient; presentation plane), and the observer was asked to transpose it onto a frontal surface (test plane). In the complementary task, a segment was shown on a frontal surface, and the observer was requested to transpose it onto an induced inclined surface. The results show that, for the first transformation, accuracy fit a linear model better, where the error increased for more inclined surfaces, that is, for larger differences between the presentation plane and the test plane. However, for the second transformation, the precision was better described by a non-linear model. The results suggest there are different mechanisms for tilt (2D orientation) and inclination processing (3D orientation).

The position of a reference matters when judging egocentric distances

R. Sousa, E. Brenner and J.B.J. Smeets (Faculty of Human Movement Sciences, Vrije Universiteit, The Netherlands).

When subjects have to point at a sphere in total darkness, with only vergence providing information about the position of the sphere, their judgments are biased. To examine whether the presence of other visible structures would reduce the bias, we examined two additional conditions. In one of the conditions, there was a reference cube at a fixed position (within each session). In the other condition, a similar cube was presented at a random position for each new sphere. We found that, regardless of whether the reference cube's position was stable, it improves the accuracy of responses. However, the improvement depended on the reference's position relative to the target. When the cube was further away from the subject than the target, the pointing responses became more accurate, but when the cube was closer to the subject than the target, it did not make any difference. It appears that subjects only use objects that are further away to calibrate distance.

Supported by FCT.

Classification of face gender and expression in different spatial frequency bands

L. Aguado (Universidad Complutense de Madrid, Spain), S. Rodríguez (Universidad Complutense de Madrid, Spain), F.J. Román (Universidad Complutense de Madrid, Spain) and I. Serrano-Pedraza (Institute of Neuroscience, Newcastle University, Newcastle upon Tyne, United Kingdom).

The role of different spatial frequency bands on gender and expression categorization was studied in three experiments. Reaction times were measured for unfiltered, low-pass (cut-off frequency of 1 cycle/deg) and high-pass (cut-off frequency of 3 cycles/deg) filtered faces. Filtered and unfiltered faces were equated in root-mean-squared contrast. For low-pass filtered faces, reaction times were higher than for unfiltered and high-pass filtered faces in both categorization tasks. In the expression task, these results were obtained with expressive faces presented in isolation (Experiment 1) and also with neutral-expressive dynamic sequences, where each expressive face was preceded by a briefly presented neutral version of the same face (Experiment 2). For high-pass filtered faces, different effects were observed on gender and expression categorization. Whereas both speed and accuracy of gender categorization were reduced in comparison to unfiltered faces, the efficiency of expression classification remained similar. Finally, we found no differences in reaction times between the effects of spatial frequency filtering on gender categorization with expressive and non-expressive faces (Experiment 3). These results show a common role of information from the high spatial frequency band in the categorization of gender and expression.

This study was supported by the project SEJ2006-01576/PSIC.

Facilitation of object recognition: An effect of contextual cueing

M. van Asselen, J. Sampaio and M. Castelo-Branco (Escola de Psicologia, Universidade do Minho, Braga, Portugal).

Implicit contextual cueing refers to a top-down mechanism in which visual search is facilitated by learned contextual features. In the current study, we aim to investigate the mechanisms underlying object-based implicit contextual cueing by studying eye movements. A visual search task was used in which participants ($n = 16$) had to locate a target object (abstract line drawing with an opening) among a number of distracters (closed abstract line drawing). In half of the trials, the set of objects presented was repeated (Repeated trials), whereas in the other trials, a new set of objects was used (New trials). The location of the objects was randomly defined, and the objects in the Repeated and New trials were repeated the same number of times. Eye movements were recorded with an iViewX High-speed Eye-Tracker. Subjects located the target faster in the Repeated trials as compared with New trials ($p < .001$), indicating that the sets of objects were memorized and search was facilitated. Shorter fixation durations were found for the Repeated trials than for the New trials [$t(20) = 3.6, p < .01$]. Furthermore, the Time from Last Fixation until Button Press (TLFBP) was significantly shorter in the Repeated than in the New trials [$t(20) = 4.6, p = .001$]. No effect was found for the Number of Fixations and the Saccade Amplitude. This indicates that object recognition is facilitated when object information is used as contextual cue.

Perceptual learning in humans: The use of same/different tasks

P.P. Faria and P.B. Albuquerque (Escola de Psicologia, Universidade do Minho, Braga, Portugal).

Perceptual learning refers to discrimination improvement between two very similar stimuli seen after non-supervised pre-exposure to these stimuli. Moreover, if two similar stimuli are pre-exposed in several alternated trials, discrimination is maximized, compared to several blocked pre-exposures of each stimulus—the intermixed/blocked effect. Recent studies in this area have used same/different tasks—in which the pre-exposed stimuli are shown in succession and participants have to classify them as being the same or different stimuli—to evaluate the effect of pre-exposure in stimulus discrimination by human subjects. Following this kind of experiments, Mitchell, Nash, and Hall (2008) propose a model of attention modulation and differential codification of features in memory as the framework for the intermixed/blocked effect. Inserted in a broader study with the objective of testing this hypothesis, the experiment presented here, using one such task, shows that: (a) the analysis of the learning that might take place during test seems to demonstrate an effect of pre-exposure in the increase of discriminability from trial to trial, regardless of pre-exposure condition; (b) overall performance in this kind of task is no different after blocked pre-exposure versus no pre-exposure; and (c) discrimination between stimuli, one pre-exposed in block and another pre-exposed intermixed, is worse than discrimination between two intermixed stimuli. Possible implications for the use of this kind of task in perceptual learning studies in humans are discussed. The work reported here was included in a Master's thesis submitted by the first author to the Universidade do Minho, Portugal.

Research supported by a grant from the Portuguese Foundation for Science and Technology (FCT) to the second author.

An exploratory study of diverse methodological factors related to the perception of beauty

P. Bustos, J. Christensen, N. Gut, A. Flexas, M. Nadal and E. Munar (Human Evolution and Cognition. IFISC (UIB-CSIC), Universitat de les Illes Balears).

Recently, there has been a great increase of interest in the neural basis of visual beauty perception. However, neuroimaging procedures differ from the usual behavioral procedures, particularly, in the use of different scales to measure responses (dichotomous vs. Likert beauty judgments). The purpose of this experiment is to explore whether any of the possible methodological options produced differences in the way 64 university students (32 men and 32 women) responded to a beauty-rating task. We analyzed their responses on Likert (1 to 9 points) and dichotomous (beauty-non beauty) scales, order (dichotomous first or Likert first), and kind of visual stimuli: artistic (impressionist, postimpressionist, abstract, realistic paintings) and non-artistic (photographs). Our results revealed that differences between sexes were only appreciable using a Likert scale, particularly in relation to the time that men and women took to decide about the beauty of realistic paintings and responses to postimpressionist paintings. On the other hand, order had an effect only on reaction times to photographs, impressionist paintings on the Likert scale, and to photographs on the dichotomous scale. However, there was no interaction between order and sex regarding reaction times and responses on either scale. Conversely, a high correlation between responses on dichotomous and Likert scales for both sexes was observed. This suggests that order of presentation of the visual stimuli has an influence on the way participants perceive beauty and that it must be taken into account in future research.

The neural bases of the perception of beauty: Differences and similarities between men and women

A. Flexas, J. Christensen, N. Gut, M. Nadal, P. Bustos and E. Munar (Human Evolution and Cognition. IFISC (UIB-CSIC), Palma de Mallorca, Spain).

Perception of beauty (aesthetic judgment) appears to be one of humans' most distinctive traits. Neuroimaging techniques are proving to be instrumental in the understanding of the biological basis of this trait. Results of recent studies suggest that a broad array of brain regions are involved in our appreciation of beauty. However, the use of different techniques makes it difficult to compare across studies. Our objective in the present work was to examine sex-related differences in the neural underpinnings of beauty appreciation by means of functional Magnetic Resonance Imaging (fMRI) and to compare these results with those of an analogous study carried out earlier by means of magnetoencephalography. Here, we recorded the BOLD signal of 12 male and 12 female participants while they decided whether diverse visual stimuli were beautiful or not beautiful. Our results suggest that there are interesting differences in specific brain regions and in lateralization. However, both in men and women, fMRI revealed occipito-temporal activity that could not be registered with magnetoencephalography due to its limitations measuring activity below cortical surface. The present results are discussed in relation to a model of beauty appreciation of visually perceived objects.

Eye movement patterns during the scanning of emotional images

R. Cabestrero (Dpto. Psicología Básica II. Facultad de Psicología. UNED), P. Quirós (Dpto. Psicología Básica II. Facultad de Psicología. UNED), A. Crespo (Dpto. Psicología Básica II. Facultad de Psicología. UNED) and P.A. Conde-Guzón (Área de Personalidad, Evaluación y Tratamiento Psicológicos. Universidad de León).

There is a close relation between eye movements and emotional experience. Hermans, Vansteenwegen, and Eelen (1999) found that during visual scanning of spider images, spider-anxious participants exhibited a different exploration patterns than controls. Isaacowitz (2005) identified different attentional preferences of optimists and pessimists toward negative emotional stimuli (looking a melanoma image). Xing and Isaacowitz (2006) reported shorter fixation durations when participants to whom emotional regulation processes were induced looked at negative emotional stimuli, in relation to positive ones. The aim of this study was to determine whether the emotional content of a stimulus induces differential gaze patterns and whether they are related to the emotional valence and activation ratings. Eye movements of 125 participants (89 women and 36 men) were recorded while looking at 16 emotional images belonging to the IAPS (International Affective Picture System). Pictures were selected taking into account the extreme ratings given to them by the Spanish sample (Moltó et al., 2001) on the orthogonal dimensions of valence and activation. Results indicated that valence dimension affected the image scanning pattern. Images with negative emotional content obtained more fixations and shorter fixation durations, as well as shorter scanning times.

Exploratory eye movements and the Müller-Lyer illusion

A. Díaz, M.P. Aivar and D.M. Jacobs (Dpto. Psicología Básica, Universidad Autónoma de Madrid, Spain).

Eye movements are often considered in explanations of visual illusions. However, in the case of the Müller-Lyer illusion, explanations based on eye movement were largely abandoned when it became apparent that the illusion also occurs under retinal stabilization (Evans & Marsden, 1966). Even so, there has been a recent renewal of interest in the analysis of the patterns of exploration that occur while observing Müller-Lyer configurations (e.g., van Doorn et al., 2009). This interest derives from Goodale and Milner's (1992) dual model, which holds that information is processed separately depending on whether it is related to conscious perception or needed for the control of action. One of the lines of evidence for this dual model is that illusions, it is argued, affect perception but not action. However, van Doorn et al. observed differences in the exploratory eye movements that underlie perception and action. This seems to imply that the differential effects of illusions on perception and action are at least partly due to the use of different informational variables, rather than only to the differential processing of information as proposed by Goodale and Milner. With the present study, we aim to provide more detail about the relation between exploratory eye movements and illusory effects. Participants observed Müller-Lyer configurations during 2 seconds, without any constraints on eye movements, before they performed a length-estimation task. In agreement with previous results, we found that the amplitude of saccades is significantly influenced by the configuration of the figure (wings-in versus wings-out). Furthermore, our results show different patterns of exploratory movements for different individuals. Some of these exploratory patterns seem to be more effective than others, because the errors in the estimation task are much smaller for some participants than for others.

What fixations reveal about perceptual styles

M.S. Puig (Vision and Control Action (VISCA-lab), Facultat de Psicologia, Universitat de Barcelona), L.P. Zapata (Vision and Control Action (VISCA-lab), Facultat de Psicologia, Universitat de Barcelona), A. Aznar-Casanova (Vision

and Control Action (VISCA-lab), Facultat de Psicologia, Universitat de Barcelona) and H. Super (Vision and Control Action (VISCA-lab), Facultat de Psicologia, Universitat de Barcelona and Institució Catalana de Recerca i Estudis Avançats (ICREA), Barcelona).

Perceptual style refers to the way we perceive information and how we use this information to solve problems. At the extremes of perceptual style scores, two styles can be distinguished: field dependence and field independence. In the latter case, observers have a strong tendency for visual detail (local) processing whereas in the former case, observers perceive more globally. Such a difference is reflected in superior performance in the embedded figures test for field independent subjects. These subjects are better at detecting a simple figure embedded in a complex shape than are field dependent subjects. So, independent observers are less influenced by the context of a visual stimulus than dependent observers. Besides perception, it is well known that the context of a stimulus also affects eye movements, that is, during visual search. Until now, there has been no evidence of whether field dependent and independent observers display different gaze patterns, and whether it relates to their discrepancy in visual performance. Therefore, in the first part of this study, we registered eye movements of observers performing an embedded figures test, and analyzed their gaze pattern in relation to the embedded figure. In the second part, we recorded eye gaze of field dependent and independent observers, as characterized by the Witkin test, during an embedded figures test. The results are discussed in terms of different visual strategies for solving the embedded figures test, and in terms of characterizing perceptual style.