

# REALIZED GENETIC PARAMETERS OF GROWTH AND REPRODUCTIVE TRAITS AFTER 25 YEARS OF SELECTION IN THE ASTURIANA DE LOS VALLES BEEF CATTLE BREED

25 AÑOS DE SELECCIÓN EN LA RAZA DE CARNE ASTURIANA DE LOS VALLES: ANÁLISIS DE LOS PARÁMETROS GENÉTICOS EN CARACTERES DE CRECIMIENTO Y REPRODUCTIVOS.

Cortes O.<sup>1</sup>, Carleos C.<sup>2</sup>, Baro J.A.<sup>3</sup>, Fernández M.A.<sup>4</sup>, Villa J.<sup>4</sup>, Menéndez-Buxadera A.<sup>1</sup>, Cañon J.<sup>1\*</sup>

<sup>1</sup>Departamento de Producción Animal. Facultad de Veterinaria. Universidad Complutense de Madrid. Madrid. España. jcanon@vet.ucm.es

<sup>2</sup>Dpto. Estadística e Investigación Operativa. Universidad de Oviedo. Oviedo. España

<sup>3</sup>Dpto. CC. Agroforestales. ETSIIAA. Universidad de Valladolid. Palencia. España

<sup>4</sup>ASEAVA. Llanera. Asturias. España

## Keywords:

Heritability  
Breeding values  
Calving ease  
Calving interval  
Weaning weight

## Palabras clave:

Heredabilidad  
Valores  
mejorantes  
Facilidad de parto  
Intervalo entre  
partos  
Peso al destete

## Abstract

After 25 years of selection, genetic parameters estimates in the Asturiana de los Valles cattle breed are revisited. Two reproduction traits, calving ease and calving interval, and one growth trait, weaning weight, were analysed for this study. A total of 198,277 records for calving ease, 51,161 for weaning weights, and 123,532 for calving interval were available. Uni- and bi-variate model were used in the analysis, and the maternal effect was included for WW. Heritabilities for reproductive traits were lower than for growth traits and the univariate and bivariate models achieved similar heritability estimates. The maternal effect was smaller than the direct genetic effect for WW, and negative correlation was detected among them in accordance with similar studies for cattle beef breeds. The genetic correlation among WW and CE was moderate and negative (-0.262) and the WW-CE genetic correlation was quite low (0.056). The bivariate model exposed an increase in the accuracy of the estimated breeding values for the three traits analyzed. Genetic trends were estimated to evaluate the genetic change over the last 25 years. The highest genetic progress was attained by WW, with a regression coefficient of the breeding values on the year estimated at 0.225. The genetic trend for CE was negligible and CI evidenced a small, negative genetic change (-0.128)

## Resumen

Se han analizado los parámetros genéticos de dos caracteres reproductivos, facilidad de parto e intervalo entre partos y un carácter de crecimiento, peso al destete, en la raza de carne Asturiana de los Valles después de 25 años de selección como una representación de los principales caracteres de interés económico en esta raza. En el análisis se utilizaron modelos uni y bivariado incluyendo el efecto materno en el peso al destete en ambos modelos. La heredabilidad estimada para el carácter de crecimiento fue superior a la de los caracteres reproductivos. Los modelos uni y bivariado dieron estimas de heredabilidad similares para todos los caracteres. En el peso al destete el efecto materno estimado fue inferior al efecto genético directo y su correlación resultó negativa, como se ha descrito previamente en otros estudios. Las correlaciones genéticas entre los caracteres fueron -0.262 entre peso al destete e intervalo entre partos y de 0.056 peso al destete y facilidad de parto. La fiabilidad de los valores genéticos estimados fue mayor con el modelo bivariado. En los 25 años de selección el mayor progreso genético corresponde con el peso al destete, con un coeficiente de regresión de 0.225, para el intervalo entre partos el progreso genético también fue favorable, con un coeficiente de regresión negativo (-0.128), y para la facilidad de parto no se observó una tendencia significativamente diferente de cero.

## Introduction

Beef cattle farming is an important component of sustainable agriculture in much of Europe. The large variety of environmental, social and marketing conditions in Europe has favored the maintenance of a wide spectrum of beef cattle breeds, especially in the South of Europe as meat consumption of local cattle breeds is higher than in

North. Periodic reviews of beef improvement programs are both for benchmarking and as an opportunity to evaluate their efficacy, to understanding the phenotypic trends and to estimate the genetic parameters changes of the traits under selection. Asturiana de los Valles is an autochthonous beef cattle breed mainly located in the north of Spain (Asturias). As in any beef cattle breed, traits related to reproduction and growth should be considered as the main variables within the breeding objectives (Phocas *et al.*, 1998; Albera *et al.*, 2004). In beef cattle, weights at different ages (birth, weaning or yearling weights) or weight over certain ages interval have been traditionally adopted as the selection criteria. Weight traits are easy to measure and have high correlations among them. Also a high correlation has been demonstrated among weight measures at early ages and calving ease (CE) (Meijering *et al.*, 1984; Koots *et al.*, 1994). So, an increase in the birth or weaning weight (WW) may be detrimental for calving ease comprising the traditionally semi-extensive rearing of the Asturiana de los Valles breed. Furthermore, this breed is regarded as well adapted to local environmental conditions and maternal behaviour could be affected as growth traits increase. The heritability estimated for growth traits, like weaning weights, is relatively high, (Koots *et al.*, 1994; Phocas & Lalöe 2004; Roughsedge *et al.*, 2005) but is lower for reproductive traits, especially in those traits, like calving ease, where phenotype scores are sensitive to subjectivity and strongly influenced by non-genetic factors (MacNeil *et al.*, 1984; Cundiff *et al.*, 1986; Meyer *et al.*, 1990). So, for reproductive traits larger data sets are required to obtain reliable genetic parameter estimates. Finally, in traits measured at an early age, phenotypes are affected by two separated divergent components, the calf effect and the maternal effect, which contribute to making the definition of breeding objectives difficult (Phoca & Lalöe 2004). In this complex situation where different traits, some with negative genetic correlations, are economically important, the estimation of genetic parameters is crucial to planning breeding programmes and to understanding the phenotypic trends of the traits under selection. There are published studies based on growth and reproductive traits in cattle breeds (Eriksson *et al.*, 2004; Cammack *et al.*, 2009; Casas *et al.*, 2010; MacNeil & Vukasinovic, 2011). However, the information available on the genetic relationship between calving ease and calving interval (CI), as reproduction traits, and weaning weight, as a growth trait, is relatively scarce. In spite of the great number of phenotypes registered in the Asturiana de los Valles breed program, previous analysis for growth and reproductive genetic parameter traits (Gutierrez *et al.*, 1997; Goyache *et al.*, 2005; Gutierrez *et al.*, 2006; Gutierrez *et al.*, 2007; Cervantes *et al.*, 2010; Pun *et al.*, 2011) have been based on relatively limited databases, including only animals registered until 1997. Therefore, those genetic parameter estimates should be jointly revisited, along with their effect on the different trait trends. The aims of the current study were (1) to estimate genetic parameters of growth and reproductive traits, taking into account the direct and maternal genetic effect and the genetic correlations between them and (2) to analyze the trend of the breeding values after 25 years' selection.

## Material and methods

The Asturiana de los Valles bovine breed is located in the north of Spain over the Cantabrian range under semi-extensive conditions. Animals graze on communal pastures during 8 to 9 months and winter on valley pastures that need supplemental hay. The initial phenotypic and pedigree database covered 24 years (1987-2011) of information on the Asturiana de los Valles beef cattle breed.

### Animals

Pedigree information on the Asturiana de los Valles breed was provided by the official breed association (ASEAVA). Those animals with identification or birth weight errors were excluded for further analysis. Finally, a total of 276,654 pedigree records were included in the statistical analysis.

### Traits

Reproductive and growth traits analyzed included weaning weight (WW), calving ease (CE) and calving interval (CI). Phenotypic records of those animals with birth or identification errors were excluded for further analysis. Also, values lower than 300 days or higher than 730 for CI days were discarded.

Traditionally, calving ease was recorded following the criteria: (1) no assistance; (2) minor assistance; (3) hard assistance; and (4) caesarean. In our data scores 1 and 2 were considered as a single group due to slight differences between them.

After editing the database, a total of 198,277 calving ease records registered in 76,895 cows, 51,161 weaning weights records registered in 25,495 cows, and 123,532 calving interval records registered in 44,812 cows were included for the analysis.

An inflection point for phenotypic records with errors was detected at 1997 and consequently, previous records were dropped and the final phenotypic database included information from 1997 to 2011. It must be noted that several reports (Gutierrez et al. 1997; Goyache et al. 2005; Gutierrez et al. 2006; Gutierrez et al. 2007; Cervantes et al. 2010; Pun et al. 2011) for growth and reproductive traits in Asturiana de los Valles cattle have been based on records registered until 1997, while such records have been rejected in our final database due to the large number of errors found. Table I shows the number of observations, the mean and standard deviations and the minimum and maximum values for each trait.

**Table I.** Number of observations, mean and standard error for weaning weight (WW) and calving interval (CI). Frequencies (in percentage) for calving ease (CE). (*Número de observaciones, media, desviación estándar para los caracteres peso al destete e intervalo entre partos. Frecuencias en porcentaje para el carácter dificultad al parto*)

Trait	N	Mean	SD
Weaning weight (WW)	51,161	219.7 <sup>1</sup>	55.2
Calving interval (CI)	123,532	418.1 <sup>2</sup>	99.2
	Calving ease (CE)		
	1 <sup>3</sup>	2 <sup>3</sup>	3 <sup>3</sup>
Frequency (%)	97.4	1.6	1

<sup>1</sup>In kilograms

<sup>2</sup>In days

<sup>3</sup>1 null or minor assistance; 2 hard assistance; 3 caesarean

### Statistical analysis

Data were analyzed using ASReml 3 (Gilmour et al. 2000). The non-genetic effects in the models included as fixed effects were: the number of calving (12 levels) and year-month of calving (156 levels); for both WW and CE the model also included the sex (2 levels), and for CE the age at weaning was also included as a cubic covariate. The random effects taken into consideration were direct and maternal genetic effects, the permanent environmental effect, and the combination management-municipality as a temporal effect for WW (324 levels); the maternal effect was excluded from the models for CI and CE.

The components of co-variance were estimated using a univariate and a bivariate model combining WW with CI and with CE. The matrix notation for the univariate model was:  $y = Xb + Za + Wm + Pp + Qq + e$

$$\begin{bmatrix} a \\ m \\ p \\ q \\ e \end{bmatrix} \approx N \left( \begin{bmatrix} 0 \\ 0 \\ 0 \\ 0 \\ 0 \end{bmatrix}, \begin{bmatrix} G & C & 0 & 0 & 0 \\ C & M & 0 & 0 & 0 \\ 0 & 0 & P & 0 & 0 \\ 0 & 0 & 0 & Q & 0 \\ 0 & 0 & 0 & 0 & R \end{bmatrix} \right)$$

where  $G = A\sigma_a^2$ ,  $M = A\sigma_m^2$ ,  $C = A\sigma_{am}$ ,  $P = Ip\sigma_p^2$ ,  $Q = Iq\sigma_q^2$ , and  $R = In\sigma_e^2$ .  $\mathbf{Y}$  is the vector of observations for WW, CE or CI;  $\mathbf{b}$  is a vector of non-genetic effects;  $\mathbf{a}$  is a vector of the random direct additive genetic effects;  $\mathbf{m}$  is a vector of the random maternal additive genetic effects;  $\mathbf{p}$  is a vector of random permanent environmental effects,  $\mathbf{q}$  is the vector of random management-municipality effects, and  $\mathbf{e}$  is the vector of residuals.  $\mathbf{X}$ ,  $\mathbf{Z}$ ,  $\mathbf{W}$ ,  $\mathbf{P}$ , and  $\mathbf{Q}$  are incidence matrices relating  $\mathbf{b}$ ,  $\mathbf{a}$ ,  $\mathbf{m}$ ,  $\mathbf{p}$ , and  $\mathbf{q}$ , to  $\mathbf{y}$ .  $\mathbf{A}$  is the additive numerator relationship matrix that is created using pedigree information.  $\sigma_a^2$  is the variance of direct additive genetic effects,  $\sigma_m^2$  is the variance of maternal additive genetic effects,  $\sigma_{am}$  is the covariance between direct and maternal additive genetic effects,  $\sigma_p^2$  and  $\sigma_q^2$  are the variances of permanent environmental and management-nucleus effects respectively, and  $\sigma_e^2$  is the residual variance.  $\mathbf{Ip}$ ,  $\mathbf{Iq}$ , and  $\mathbf{In}$  are identity matrices with order respectively equal to  $\mathbf{p}$ , the number of environments;  $\mathbf{q}$ , the number of management-municipality levels; and  $\mathbf{n}$ , the number of observations. In the univariate models for CE and for CI the random maternal additive genetic effects ( $\mathbf{m}$ ) and the covariance between direct and maternal genetic effects (C) were excluded.

Additive direct heritability ( $h_{ai}^2$ ) and additive maternal heritability ( $h_{mi}^2$ ) were estimated as ratios of additive direct and additive maternal variances to phenotypic variance, respectively. The direct-maternal genetic

correlation ( $r_{a,m}$ ) was computed as the ratio of the direct-maternal genetic covariance ( $\sigma_{a,m}$ ) to the product of the square roots of  $\sigma_a^2$  and  $\sigma_m^2$ .

Traits WW-CE and WW-CI were analysed following the model:

$$\begin{bmatrix} y_1 \\ y_2 \end{bmatrix} = \begin{bmatrix} X_1 & 0 \\ 0 & X_2 \end{bmatrix} \begin{bmatrix} b_1 \\ b_2 \end{bmatrix} + \begin{bmatrix} Z_1 & 0 \\ 0 & Z_2 \end{bmatrix} \begin{bmatrix} a_1 \\ a_2 \end{bmatrix} + \begin{bmatrix} W_1 & 0 \\ 0 & W_2 \end{bmatrix} \begin{bmatrix} m_1 \\ m_2 \end{bmatrix} \\ + \begin{bmatrix} P_1 & 0 \\ 0 & P_2 \end{bmatrix} \begin{bmatrix} p_1 \\ p_2 \end{bmatrix} + \begin{bmatrix} Q_1 & 0 \\ 0 & Q_2 \end{bmatrix} \begin{bmatrix} q_1 \\ q_2 \end{bmatrix} + \begin{bmatrix} e_1 & e_{21} \\ e_{12} & e_2 \end{bmatrix}$$

In this model, vectors and incidence matrices correspond to those in the univariate model, and subscripts 1 and 2 denote traits. For example,  $\mathbf{a}$  is now a vector resulting from concatenating  $\mathbf{a1}$  and  $\mathbf{a2}$ . The changing elements of the formula are now the following:

$G = G_0 \otimes A$ ,  $M = M_0 \otimes A$ ,  $C = C_0 \otimes A$ ,  $P = Ip\sigma_p^2$ ,  $Q = Iq\sigma_q^2$ , and  $R = In\sigma_e^2$ .  $G_0$  is a 2x2 covariance matrix between direct additive genetic effects for traits 1 and 2.  $M_0$  is a 2x2 covariance matrix between maternal additive genetic effects for traits 1 and 2.  $C_0$  is a 2x2 covariance matrix between direct additive genetic effects for traits 1 and 2. So we have:

$$var \begin{pmatrix} a_1 \\ a_2 \\ m_1 \\ m_2 \end{pmatrix} = \begin{bmatrix} G & C \\ C & M \end{bmatrix} = \begin{bmatrix} \sigma_{a_1}^2 & \sigma_{a_1 a_2} & \sigma_{a_1 m_1} & \sigma_{a_1 m_2} \\ \sigma_{a_1 a_2} & \sigma_{a_2}^2 & \sigma_{a_2 m_1} & \sigma_{a_2 m_2} \\ \sigma_{a_1 m_1} & \sigma_{a_2 m_1} & \sigma_{m_1}^2 & \sigma_{m_1 m_2} \\ \sigma_{a_1 m_2} & \sigma_{a_2 m_2} & \sigma_{m_1 m_2} & \sigma_{m_2}^2 \end{bmatrix} \otimes A$$

The correlations were estimated for direct genetic effect among the univariate and bivariate models for WW, CE and CI and for maternal genetic effect for WW. Genetic trends were calculated as the regression of the average predicted breeding values of the animals born in the same year on year. The following equation was used:

$$y_i = b_0 + b_1 x_i + e_i$$

in which  $y_i$  is the mean breeding value for the traits evaluated in the  $i$ th birth year,  $b_0$  is the intercept,  $b_1$  is the slope,  $x_i$  is the  $i$ th birth year, and  $e_i$  is a random error. Regression coefficients and goodness of fit ( $R^2$ ) values were also derived as part of the procedure.

The accuracy of the breeding values was estimated using:

$$r_{\hat{H}H} = \sqrt{1 - \frac{\sigma_e^2}{\sigma_{ai}^2}} \times 100$$

Where  $\sigma_e^2$  denotes the prediction error variance for each animal and trait, and  $\sigma_{ai}^2$  is the additive genetic variance for each trait analyzed.

**Results**

Phenotypic variances and genetic parameter estimates in the univariate model are given in Table II. The additive variance component relative to the phenotypic one was greater for WW than for CE and CI (0.506, 0.088 and 0.227, respectively). Maternal heritability (0.218) was lower than that in direct genetic effect for WW. The genetic correlation for direct-maternal direct genetic effect was negative for WW (-0.632).

Bivariate genetic parameter estimates and their correlations are shown in Table III. The direct (WW, CE and CI) and maternal heritabilities (WW) were similar to those in the univariate model. Also, the correlation between direct and maternal effect for WW was negative and evidenced similar values in bivariate and univariate models (Table II and 3). The genetic correlation for direct genetic effect in the bivariate model was positive and low in WW-CE (0.056) and negative and higher in the WW-CI bivariate model (-0.262).

The genetic correlations among the total breeding values obtained by the univariate and the bivariate models are shown in Table IV. The high genetic correlation between models for the total breeding values estimated is remarkable. The average of the breeding value accuracies of all the animals analyzed and of the sires with more than 15 offspring was higher in the bivariate than in the univariate model.

The evolution of the breeding values from 1997 to 2011 was estimated from the regression coefficient of the breeding values on the birth (Table V). While CE and CI showed null and negative regression values (-0.0004 and -0.128, respectively), the regression coefficient for WW was positive (0.267).

**Table II.** Estimates of phenotypic variances, direct and maternal heritabilities, total heritability and correlations between direct and maternal genetic effects for all traits analysed in the univariate model. Standard errors are in parentheses (*Estimas del modelo univariado de las varianzas fenotípicas, heredabilidad directa y materna, heredabilidad total y las correlaciones entre el efecto genético directo y materno para los caracteres analizados. Errores estándar entre paréntesis*)

	CE	WW	CI
Animals, No	246,546	89,925	164,532
Animals with records, No	198,277	51,161	123,532
Direct genetics	4934*	888.66	2312.91
Maternal genetics		382.31	
Covariance direct-maternal		-368.66	
Permanent environment	1442.4*	10.039	235.79
System-municipality	1248*	169.85	155.07
Residual	49504*	672.73	7495.07
Total variance	56005.2*	1754.9	10198.84
$h_a^2$	0.088 $\pm$ 0.004	0.506 $\pm$ 0.021	0.227 $\pm$ 0.004
$h_m^2$		0.218 $\pm$ 0.017	
$r_{g[a,m]}$		-0.632 $\pm$ 0.025	

\*x10<sup>6</sup>

**Table III.** Estimates of phenotypic variances, direct and maternal heritabilities, total heritability, correlations between direct and maternal genetic effects and correlation between CI/WW and CE/WW following the bivariate model. Standard errors are in parentheses (*Estimas del modelo bivariado de las varianzas fenotípicas, heredabilidad directa y materna, heredabilidad total y las correlaciones entre el efecto genético directo y materno y entre el intervalo entre partos/peso al destete y dificultad al parto /peso al destete. Errores estándar entre paréntesis*)

	Traits bivariate model		Traits bivariate model	
	CE	WW	CI	WW
Animals, No	246,546		164,532	
Animals with records, No	198,277		123,532	
Direct genetics	4918*	941.6	2406.3	970,4
Maternal genetics	-	376.9	-	306,9
Covariance direct-maternal	-	-417.3	-	-375,7
Permanent environment	1441*	254.605	172.981	585.954
System-municipality	124*	166.110	144.560	148.274
Residual	49513*	677.025	7505.84	605.463
Total variance	55390*	1769.7	9610.8	1713.9
$h_a^2$	0.089 (0.004)	0.532 (0.022)	0.25 (0.008)	0.566 (0.026)
$h_m^2$	-	0.213 (0.018)	-	0.179 (0.026)
$r_{g[a,m]}$	-	-0.700 (0.023)	-	-0.688 (0.034)
$r_a$	0.056 (0.040)		-0.262 (0.037)	

\*values x10<sup>6</sup>

**Table IV.** Breeding value correlations between univariate and bivariate models for calving ease (CE), calving interval (CI) and weaning weight (WW). Accuracies of the estimated breeding values in all the animals analyzed and sires with more than 15 offspring in the univariate and the bivariate models (*Correlaciones de los valores genéticos estimados con el modelouni y bivariado en los tres caracteres analizados, peso al destete, dificultad al parto e intervalo entre partos. Fiabilidad de los valores genéticos estimados en todos los animales analizados y en los sementales con más de 15 descendientes*)

Breeding value correlations	CI	CE	WW			
EBV animals	0.968	0.996	0.996			
EBV sires	0.993	0.996	0.986			
Breeding value accuracies	Univariate	Bivariate	Univariate	Bivariate	Univariate	Bivariate
Accuracy animals	29.6	40.8	37.4	45.3	41.1	53.5
Accuracy sires	39.1	64.8	54.5	65.5	51.4	66.3

**Table V.** Regression coefficients of the breeding values on year and goodness of fit ( $R^2$ ) for the three traits analysed, Calving ease (CE), calving interval (CI) and weaning weight (WW) (*Coefficiente de regresión de los valores genéticos sobre el año y bondad del ajuste ( $R^2$ ) en los tres caracteres analizados, dificultad al parto, intervalo entre partos y peso al destete*)

Trait	Regression coefficient	$R^2$ (%)
CE	-0.0004	80
CI	-0.128*	82
WW	+0.225*	74

\* $p < 0.05$

## Discussion

Traditionally the selection criteria of the Asturiana de los Valles farmers have been focused on morphology, well conformed animals, and in the double-muscled character more than in breeding values for growth or reproductive traits. Currently, the frequency of the myostatin (*mh*) allele is 82% and the *mh* homozygote (*mh/mh*), heterozygote (*mh/+*) and normal (*+/+*) genotype frequencies are 69%, 27% and 4%, respectively (data not shown). The *mh* allele has a significant effect on growth traits, a *mh* homozygote calf weight 5 kg more than a normal animal at birth but between 4 and 10 kg less at weaning weight (data not shown). In spite of the higher birth weight and the higher frequency of *mh* homozygote animals in the Asturiana de los Valles breed, only 2.7% of calvings required hard assistance or a caesarean. The main traits included in the beef cattle genetic programs are related to growth traits; however, the relative importance of reproduction traits could be up to 4-fold more important than improvements in production traits when calves are sold at weaning (Melton, 1995). In general, reproductive traits show low heritabilities and are recorded later in the life of the animal than the majority of growth traits, so their improvement has been mainly carried out through crossbreeding and improved management techniques rather than direct selection. However, the important genetic correlations between reproductive traits and others traits with moderate or high heritabilities made indirect selection for reproductive traits feasible (Camamck et al. 2009).

As expected, the results showed that direct genetic heritabilities for reproductive traits (CE and CI) were lower than for production traits (WW). While fertility is a general term and not easily defined, CE has been easily defined and categorized. Notwithstanding, heritabilities for CE were low and of similar magnitude in the univariate and bivariate models ( $\sim 0.09$ ), suggesting that direct selection for CE will result in a low response in an improvement programme of reduced size. Also, the direct genetic heritability for CE (0.09) was lower than that in other bovine beef breeds (Koots et al. 1994; Phocas & Laloe, 2004), even lower than that in previous Asturiana de los Valles breed analyses (Gutierrez et al. 2007; Cervantes et al. 2010). In contrast the heritabilities for WW and CI, and the magnitude of the correlation among direct and maternal genetic for WW were higher than those in previous analyses in the Asturiana de los Valles and Asturiana de la Montaña breeds (Gutierrez et al. 2002; Gutierrez et al. 2006; Gutierrez et al. 2007; Pun et al. 2011; Baro et al. 2012). The differences with previous analyses in the Asturianan de los Valles breed could be due to two reasons. (1) Previous heritability estimates referred to before were based on animals born until 1997 while animals born

before this year were rejected in our database phenotypic records. (2) Also, when the covariance among direct and maternal effect is not negligible, the direct genetic effect is greater (Meyer, 1997), as recorded for CE in Cervantes et al. (2010).

As a consequence of the low CE heritability the regression coefficient for CE showed no genetic change ( $b = -0.0004$ ) from 1997 to 2011. So, taking into account the high frequency of calving ease belonging to level 1 and the low heritability value, the improvement in reproductive traits should focus more on other traits than CE and the improvement in management techniques would be more effective than direct selection in this trait. Furthermore, the current estimate of the genetic correlation between CE and direct WW (0.056) would suggest that an increase in WW is not expected to have an important effect on the correlated response of CE.

CI measure would be an indicator of reproductive health throughout the life of a cow. The direct genetic heritabilities for CI in the univariate and bivariate models ( $\sim 0.24$ ) was higher than that in previous studies in the Asturiana de los Valles breed (Goyache et al. 2002; Gutierrez et al. 2007;) and than the mean value of 0.10 reported by Koots et al. (1994). Also, Gutierrez et al. (2007) reported a negligible correlation between CI and WW ( $-0.068$ ), suggested by the high estimated SE which includes zero (0.112). In our results the genetic correlation between CI and WW showed a consistent negative value ( $-0.262$ ). CI needs two calves to be scored, so CI scores have mostly been obtained from well conformed cows that are priority selected by farmers and receive more favourable treatments from farmers and more opportunities to conceive. So a calf born from a best breed conformation cow, which should have a shorter CI, is expected to have a higher weaning weight. However, the negative correlation among CI and WW could be biased because the best conformed cows were managed differently as regards reproduction.

The genetic trend from 1997 to 2011 for CI was negative in the desired direction, but weak ( $b = -0.128$  days). It is remarkable that breeding values for CI were not available for breeders, so phenotypic values are the only source of information and, taking into account its low heritability, a weak increase in the genetic merit for this trait is to be expected if selection is based on phenotypic data. Also, short CI could be associated with cows whose first calves are born late and, in these situations, selection of cows with short CI would indirectly increase the age at first calving (Cammarck, 2009).

WW showed the higher direct genetic heritability when both univariate and bivariate models ( $\sim 0.5$ ) were adjusted. This heritability is higher than that in other bovine breeds (Robinson et al. 1996; Miller & Wilton, 1999; Montaldo & Kinghorn, 2003) and in the upper range of estimates by Groenenveld et al. (1998). The high negative correlation between direct and maternal heritability for WW indicated unfavourable interference among them. So, selection for direct effect decreased maternal ability and therefore direct and maternal effect should be considered for WW selection. This negative genetic correlation is commonly found among direct and maternal genetic effects (Groenenveld et al. 1998; Montaldo & Kinghorn, 2003; Eriksson et al. 2004). Nevertheless, Groenenveld et al. (1998) mentioned that negative correlation between direct and maternal effect is not clear. Several reasons have been suggested to explain this antagonism: a negative dam-offspring correlation due to the decrease in cow milk production when calf weaning weights increase or the adverse effect of the heifers' nutrition on the WW of their calves could explain it (El-Saied et al. 2006). Furthermore, Wilson & Reale (2006) mentioned that the negative correlation is the consequence of a pleiotropic effect in order to preserve the genetic variation that limits the selection response.

The genetic trend for WW ( $b = 0.268$  kg.) showed a slight genetic gain from 1997 to 2011. Furthermore, the phenotypic increase for WW over the last 25 years was low (7%) (data not shown) in spite of the high heritability estimated (0.53). This result suggests that the genetic progress for WW was lower than expected, probably due to the high frequency increases in mh homozygote animals (67% mh/mh) in the Asturianan de los Valles breed and their weaning weights being lower than that of normal animals. The negative correlation between the direct and the maternal genetic effect could slow down the WW phenotypic increase, as reflected the expression in Willham (1972) for selection response,  $(\sigma_a^2 + 1.5\sigma_{am} + 0.5\sigma_m^2)/\sigma_p$ , where negative  $\sigma_{a,m}$  values result in decreased responses. In order to combine both contrary genetic effects, the maternal and direct WW effects could be included in the selection criteria for the Asturiana de los Valles breed. Also, the lower offspring from artificial insemination rather than natural services sires and other criteria than breeding values used to select sires by the breeders could explain this result.

The uni- and bivariate models produced similar genetic parameter estimates and, as expected (Kadarmideen et al. 2003), high correlations between univariate and bivariate estimated breeding values for the three traits analyzed. The higher number of phenotypic observations for CI than for WW allows us to evaluate 3-fold more

animals for WW in the bivariate model (CI-WW) than in the univariate model (WW). As expected, the values of accuracies of the estimated breeding values were higher in the bivariate model than in the univariate one and higher in sires than in all the animals. Also, the higher genetic correlation between CI-WW than between CE-WW is likely to explain the higher increase in accuracy for CI than for CE in the bivariate model. So the bivariate model could be considered as an alternative method to estimate breeding values for the three traits analysed.

### Conclusions

As expected, heritability for WW was clearly higher than for CI and CE. In spite of the high heritability estimated for WW its genetic trend over the 25 years of selection was low. This is also the situation for traits CE and CI. Customary selection, as carried out by Asturiana de los Valles breeders, prioritizes morphology and double-muscled phenotypes, jointly with the pleiotropic effect of the mh allele on weaning weight, and could slow down the WW genetic trend. Our results indicated low genetic correlation between WW and CE, and moderate negative ones between WW and CI. The high negative genetic correlation estimated between direct and maternal genetic effect for WW suggests that both effects are suitable for inclusion as selection criteria for the Asturiana de los Valles beef cattle breed.

### Acknowledgements

The project has been partially funded by Ministerio de Agricultura, Alimentación y Medio Ambiente of Spain (MAGRAMA) and by the breeder association ASEAVA who also provided the information used in this study.

### References

- Albera A., Carnier P. & Groen A.F. 2004. Definition of a breeding goal for Piemontese breed: Economic and biological values and their sensitivity to production circumstances. *Livestock Production Science*, 89, 67–78.
- Baro J.A., Carleos C., Menendez-Buxadera A., Rodriguez-Castañon A. & Cañon J. 2012. Genetic variability underlying maternal traits of Asturiana de la Montaña beef cattle. *Spanish Journal of Agricultural Research*, 10, 69-73.
- Cammack K.M., Thomas M.G. & Enns R.M. 2009. Review: Reproductive Traits and Their Heritabilities in Beef Cattle. *The Professional Animal Scientist*, 25, 517–528
- Casas E., Thallman R.M. & Cundiff L.V. 2010. Birth and weaning traits in crossbred cattle from Hereford, Angus, Brahman, Boran, Tuli, and Belgian Blue sires. *Journal of Animal Science*, 89, 979-987.
- Cervantes I., Gutierrez J.P., Fernandez I. & Goyache F. 2010. Genetic relationships among calving ease, gestation length, and calf survival to weaning in the Asturiana de los Valles beef cattle breed. *Journal of Animal Science* 88, 96–101
- Cundiff L.V., MacNeil M.D., Gregory K.E. & Koch R.M. 1986. Between and within-breed genetic analysis of calving traits and survival to weaning in beef cattle. *Journal of Animal Science*, 63, 27.
- El-Saied U.M., de la Fuente L.M., Rodríguez R. & San Primitivo F. 2006. Genetic parameter estimates for birth and weaning weights, pre-weaning daily weight gain and three type traits for Charolais beef cattle in Spain. *Spanish Journal of Agricultural Research*, 4, 146-155
- Eriksson S., Näsholm A., Johansson K. & Philipsson J. 2004. Genetic relationships between calving and carcass traits for Charolais and Hereford cattle in Sweden. *Journal of Animal Science*, 82, 2269-2276.
- Gilmour A.R., Cullis B.R. & Welham S.J., Thompson R. 2000. ASREML Reference Manual. NSW Agric Biom Bull NSW Agriculture, Locked Bag, Orange, Australia.
- Goyache F., Gutierrez J.P., Fernandez I., Royo L.J. & Alvarez I. 2005. Genetic analysis of days open in beef cattle. *Livestock Production Science*, 93, 283–289.
- Gregory K.E., Cundiff L.V. & Koch R.M. 1982. Comparison of crossbreeding systems and breeding stocks used in suckling herds of continental and temperate areas. *Proc. 2rid World Congress on Genetics Applied to Livestock Production II*: 482.
- Groeneveld E., Mostert B.E. & Rust T. 1998. The covariance structure of growth traits in the Afrikaner beef population. *Livestock Production Science*, 55, 99-107.

- Gutierrez J.P., Canon J. & Goyache F. 1997. Estimation of direct and maternal genetic parameters for preweaning traits in the Asturiana de los Valles beef cattle breed through animal and sire models. *Journal of Animal Breeding and Genetics*, 114, 261–266.
- Gutiérrez J.P., Alvarez I., Fernandez I., Royo L.J., Diez J. & Goyache F. 2002. Genetic relationships between calving date, calving interval, age at first calving and type traits in beef cattle. 2002. *Livestock Production Science*, 78, 215-222.
- Gutierrez J.P., Fernandez I., Alvarez I., Royo L.J. & Goyache F. 2006. Sire x contemporary group interactions for birth weight and preweaning growth traits in the Asturiana de los Valles beef cattle breed. *Livestock Science*, 99,61–68.
- Gutierrez J.P., Goyache F., Fernandez I., Alvarez I. & Royo L.J. 2007. Genetic relationships among calving ease, calving interval, birth weight, and weaning weight in the Asturiana de los Valles beef cattle breed. *Journal of Animal Science*, 85, 69–75.
- Kadarmideen H.N. & Wegmann S. 2003. Genetic parameters for body condition score and its relationship with type and production traits in Swiss Holsteins. *Journal of Dairy Science* 86, 3685-93
- Koots K.R., Gibson J.P. & Wilton J.W. 1994. Analyses of published genetic parameter estimates for beef production traits. 1. Heritability. *Animal Breeding*, 62, 309-338.
- MacNeil M.D., Cundiff L.V., Dinkel C.A. & Koch R.M. 1984. Genetic correlations among sex-limited traits in beef cattle. *Journal of Animal Science*, 58, 1171.
- MacNeil M.D., & Vukasinovic N. 2011. A prototype national cattle evaluation for sustained reproductive success in Hereford cattle. *Journal of Animal Science* 89:1712-1718.
- Marshall D.M. 1994. Breed Differences and Genetic Parameters for Body Composition Traits in Beef Cattle. *Journal of Animal Science* 72:2745-2755.
- Meijering A. 1984. Dystocia and stillbirth in cattle. A review of causes, relations and implications. *Livestock Production Science* 11:143–177.
- Melton B.E. 1995. Conception to consumption: The economics of genetic improvement. Proc. 27th Beef Improv. Fed. Res. Symp. Annu. Meet., Sheridan, WY. Beef Improv. Fed., Manhattan, KS. p: 40.
- Meyer K., Hammond K., Parnell P.F., Mackinnon M.J. & Sivarajasingam S. 1990. Estimates of heritability and repeatability for reproductive traits in Australian beef cattle. *Livestock Production Science* 25:15.
- Meyer K. 1997. Estimates of genetic parameters for weaning weight of beef cattle accounting for direct-maternal environmental covariances. *Livestock Production Science* 52:187–199.
- Miller S.P. & Wilton J.W. 1999. Genetic relationships among direct and maternal components of milk yield and maternal weaning gain in a multibreed beef herd. *Journal of Animal Science* 77: 1155-1161.
- Montaldo H.H. & Kinghorn B.P. 2003. Additive and non-additive, direct and maternal genetic effects for growth traits in a multibreed population of beef cattle. *Archivos de Medicina Veterinaria*, 35, 243-248.
- Phocas F., Bloch C., Chapelle P., Becherel F., Renand G. & Menissier F. 1998. Developing a breeding objective for a French purebred beef cattle selection program. *Livestock Production Science*, 57, 49–65.
- Phocas F. & Lalöe D. 2004. Genetic parameters for birth and weaning traits in French specialized beef cattle breeds. *Livestock Production Science*, 89, 121–128.
- Pun A., Goyache F., Cervantes I. & Gutierrez J.P. 2011. Cytoplasmic line effects for birthweight and preweaning growth traits in the Asturiana de los Valles beef cattle breed. *Livestock Science*, 143, 177-183.
- Riley D.G., Chase C.C.Jr, Coleman S.W., Olson T.A. & Randel R.D. 2010. Evaluation of tropically adapted straightbred and crossbred beef cattle: heifer age and size at first conception and characteristics of their first calves. *Journal of Animal Science*, 88, 3173-3182
- Robinson D.L. 1996. Estimation and interpretation of direct and maternal genetic parameters for weights of Australian Angus cattle. *Livestock Production Science*, 45, 1-11.
- Roughsedge T., Amer P.R., Thompson R. & Simm G. 2005. Genetic parameters for a maternal breeding goal in beef production. *Journal of Animal Science*, 83, 2319–2329.
- Willham R.L. 1972. The role of maternal effects in animal breeding. III. Biometrical aspects of maternal effects in animals. *Journal of Animal Science*, 35, 1288-1293.
- Wilson A.J. & Reale D. 2006. Ontogeny of additive and maternal genetic effects: lessons from domestic mammals. *Am Nat* 167:E23-38. Epub 2005 Nov 29.