

## SEMINARIO

# ATOMIC-SCALE CONTROL OF GRAPHENE MAGNETISM USING HYDROGEN ATOMS

## Iván Brihuega

Departamento de Física de la Materia Condensada, Universidad Autónoma de Madrid, España  
Condensed Matter Physics Center (IFIMAC), Universidad Autónoma de Madrid, España.

**LUNES 6 DE JUNIO A LAS 12:00**

Departamento de Física de Materiales, Sala de Seminarios, UCM

Incorporating magnetism to the long list of graphene capabilities has been pursued since its first isolation in 2004. In this talk I will show how we use a scanning tunneling microscope to explore and manipulate graphene p-magnetism at an atomic level. Our work shows how the absorption of single H atoms on graphene magnetizes the graphene regions around them. In contrast to common magnetic materials, where the magnetic moments are localized in a few angstroms, the induced graphene magnetic moments extend over several nanometers and present an atomically modulated spin texture. Our measurements also prove that the induced magnetic moments couple strongly at very long distances following a particular rule: magnetic moments sum-up or neutralize critically depending on the relative H-H adsorption sites [1].




Fig. 1. Left, STM topography of a single H atom chemisorbed in graphene. Right, STS measurements of the LDOS induced by the H atom, showing the appearance of a fully polarized peak at  $E_F$ , and of graphene.

### References

- [1] H. González-Herrero, J. M. Gómez-Rodríguez, P. Mallet, M. Moaied, J. J. Palacios, C. Salgado, M.M. Ugeda, J. Y. Veuillen, F. Ynduráin, I. Brihuega, **Science**, 352, 437 (2016).