

# SADDLE-SHAPED SOLUTION TO THE INTEGRO-DIFFERENTIAL ALLEN-CAHN EQUATION

Juan Carlos Felipe-Navarro

(joint work with Tomás Sanz-Perela)



Universitat Politècnica de Catalunya - Barcelona Graduate School of Mathematics

Second BYMAT Conference


Madrid, 20-24 May 2019

# The Allen-Cahn equation

We consider

$$-\Delta u = f(u) \quad \text{in } \mathbb{R}^n$$

where  $f$  is of bistable type.



The most classical example of nonlinearity is  $f(u) = u - u^3$ .

## Conjecture 1 [De Giorgi, 1978]

Let  $u \in C^2(\mathbb{R}^n)$  be a solution of the **Allen-Cahn equation**

$$-\Delta u = u - u^3 \quad \text{in } \mathbb{R}^n$$

such that

$$|u| \leq 1 \quad \text{and} \quad \partial_{x_n} u > 0$$

in the whole  $\mathbb{R}^n$ . Then, all level sets  $\{u = \lambda\}$  of  $u$  are hyperplanes, at least if  $n \leq 8$ . Equivalently,  $u$  is a **1D solution**, that is, a function depending only on one Euclidean variable.

## Conjecture 1 [De Giorgi, 1978]

Let  $u \in C^2(\mathbb{R}^n)$  be a solution of the **Allen-Cahn equation**

$$-\Delta u = u - u^3 \quad \text{in } \mathbb{R}^n$$

such that

$$|u| \leq 1 \quad \text{and} \quad \partial_{x_n} u > 0$$

in the whole  $\mathbb{R}^n$ . Then, all level sets  $\{u = \lambda\}$  of  $u$  are hyperplanes, at least if  $n \leq 8$ . Equivalently,  $u$  is a **1D solution**, that is, a function depending only on one Euclidean variable.

- $n = 2$  [Ghoussoub & Gui, 1998]
- $n = 3$  [Ambrosio & Cabré, 2000]
- $4 \leq n \leq 8$  [Savin, 2009] with  $\lim_{x_n \rightarrow \pm\infty} u(x', x_n) = \pm 1$
- $n \geq 9$  Counterexample [del Pino, Kowalczyk & Wei, 2011]

## Theorem 2 [Jerison and Monneau, 2004]

A counterexample for the Conjecture of De Giorgi in  $\mathbb{R}^{n+1}$  can be constructed from a bounded, even with respect to each coordinate, global minimizer of

$$-\Delta u = f(u) \quad \text{in } \mathbb{R}^n.$$

## Theorem 2 [Jerison and Monneau, 2004]

A counterexample for the Conjecture of De Giorgi in  $\mathbb{R}^{n+1}$  can be constructed from a bounded, even with respect to each coordinate, global minimizer of

$$-\Delta u = f(u) \quad \text{in } \mathbb{R}^n.$$

## Theorem 3 [Modica and Mortola, 1977]

The energy functional of a rescaled version of the Allen-Cahn equation  $\Gamma$ -converges to the perimeter functional.

## Theorem 4 [Bombieri, De Giorgi and Giusti, 1969]


If  $2m \geq 8$ , the Simons cone  $\mathcal{C} = \{(x', x'') \in \mathbb{R}^m \times \mathbb{R}^m \mid |x'| = |x''|\}$  is a global minimizer of the perimeter functional.

# Saddle-shaped solutions

## Definition 5

Let  $u$  be a bonded solution of the Allen-Cahn equation in  $\mathbb{R}^{2m}$ . We say the it is a **saddle-shaped solution** if

- It is doubly radial:  $u = u(|x'|, |x''|)$ .
- It is odd with respect to Simons Cone:  $u(|x'|, |x''|) = -u(|x''|, |x'|)$ .
- It is positive in  $\mathcal{O} = \{(x', x'') \in \mathbb{R}^m \times \mathbb{R}^m \text{ st } |x'| > |x''|\}$ .



# Known results

Existence	$2m = 2$	[Dang, Fife, Peletier, '92]
	$2m \geq 4$	[Cabr�, Terra, '09]
Asymptotic behavior	$2m = 2$	[Dang, Fife, Peletier, '92]
	$2m \geq 4$	[Cabr�, Terra, '10]
Uniqueness	$2m = 2$	[Dang, Fife, Peletier, '92]
	$2m \geq 4$	[Cabr�, '12]
Monotonicity properties	$2m = 2$	[Dang, Fife, Peletier, '92]
	$2m \geq 4$	[Cabr�, Terra, '10] [Cabr�, '12]
Unstability	$2m = 2$	[Shatzman, '95]
	$2m = 4, 6$	[Cabr�, Terra, '09, '10]
Stability	$2m \geq 14$	[Cabr�, '12]
Minimality	$2m \geq 8$	

# Known results

Existence	$2m = 2$	[Dang, Fife, Peletier, '92]
	$2m \geq 4$	[Cabr�, Terra, '09]
Asymptotic behavior	$2m = 2$	[Dang, Fife, Peletier, '92]
	$2m \geq 4$	[Cabr�, Terra, '10]
Uniqueness	$2m = 2$	[Dang, Fife, Peletier, '92]
	$2m \geq 4$	[Cabr�, '12]
Monotonicity properties	$2m = 2$	[Dang, Fife, Peletier, '92]
	$2m \geq 4$	[Cabr�, Terra, '10] [Cabr�, '12]
Unstability	$2m = 2$	[Shatzman, '95]
	$2m = 4, 6$	[Cabr�, Terra, '09, '10]
Stability	$2m \geq 14$	[Cabr�, '12]
Minimality	$2m \geq 8$	?

# The nonlocal problem

Study **saddle-shaped solution** to nonlocal Allen-Cahn equation

$$L_K u = f(u) \quad \text{in } \mathbb{R}^{2m}$$

with

$$L_K u(x) = \int_{\mathbb{R}^n} \{u(x) - u(y)\} K(x, y) \, dy.$$

- Translation invariant:

$$K(x, y) = K(x - y).$$

- Rotation invariant:

$$K(z) = K(|z|).$$

- Uniformly elliptic:

$$\frac{\lambda}{|z|^{n+2s}} \leq K(z) \leq \frac{\Lambda}{|z|^{n+2s}}, \quad \text{with } 0 < \lambda \leq \Lambda.$$

# The fractional Laplacian

**Fractional Laplacian:** canonical example of integro-differential operator.

$$K(z) = c_{n,s} |z|^{-n-2s}$$

It corresponds to radially symmetric Lévy process of order  $2s$ .

**Theorem 6: Local extension problem [Caffarelli and Silvestre, 2007]**

Let  $u : \mathbb{R}^n \rightarrow \mathbb{R}$  be a smooth bounded function, we can recover its fractional Laplacian as:

$$(-\Delta)^s u = c_{n,s} \int_{\mathbb{R}^n} \frac{u(x) - u(x+z)}{|z|^{n+2s}} dz = -d_s \lim_{y \rightarrow 0} y^{1-2s} U_y(x, y),$$

where  $U$  is the solution of the problem

$$\begin{cases} \operatorname{div}(y^{1-2s} \nabla U) = 0, & \text{in } \mathbb{R}_+^{n+1}, \\ U(x, 0) = u(x), & \text{in } \partial\mathbb{R}_+^{n+1} = \mathbb{R}^n. \end{cases}$$

# Known results

		Classical	Fractional
Existence	$2m = 2$	[Dang, Fife, Peletier, '92]	[Cinti, '11, '17]
	$2m \geq 4$	[Cabr�, Terra, '09]	
Asymptotic behavior	$2m = 2$	[Dang, Fife, Peletier, '92]	[Cinti, '11, '17]
	$2m \geq 4$	[Cabr�, Terra, '10]	
Uniqueness	$2m = 2$	[Dang, Fife, Peletier, '92]	
	$2m \geq 4$	[Cabr�, '12]	
Monotonicity properties	$2m = 2$	[Dang, Fife, Peletier, '92]	[Cinti, '11, '17]
	$2m \geq 4$	[Cabr�, Terra, '10] [Cabr�, '12]	
Unstability	$2m = 2$	[Shatzman, '95]	[Cabr�, Sol�-Morales, '05]
	$2m = 4, 6$	[Cabr�, Terra, '09, '10]	[Cinti, '11, '17]
Stability	$2m \geq 14$	[Cabr�, '12]	
Minimality	$2m \geq 8$	?	

# New results

		Classical	Fractional
Existence	$2m = 2$	[Dang, Fife, Peletier, '92]	[Cinti, '11, '17]
	$2m \geq 4$	[Cabr�, Terra, '09]	
Asymptotic behavior	$2m = 2$	[Dang, Fife, Peletier, '92]	[Cinti, '11, '17]
	$2m \geq 4$	[Cabr�, Terra, '10]	
Uniqueness	$2m = 2$	[Dang, Fife, Peletier, '92]	[F-N, Sanz-Perela, '18]
	$2m \geq 4$	[Cabr�, '12]	
Monotonicity properties	$2m = 2$	[Dang, Fife, Peletier, '92]	[Cinti, '11, '17] [F-N, Sanz-Perela, '18]
	$2m \geq 4$	[Cabr�, Terra, '10] [Cabr�, '12]	
Unstability	$2m = 2$	[Shatzman, '95]	[Cabr�, Sol�-Morales, '05]
	$2m = 4, 6$	[Cabr�, Terra, '09, '10]	[Cinti, '11, '17]
Stability	$2m \geq 14$	[Cabr�, '12]	[F-N, Sanz-Perela, '18]
Minimality	$2m \geq 8$	?	?

An important consequence of our stability result:

Corollary 7: [F-N and Sanz-Perela, 2018]

The Simons cone is a stable  $2s$ -minimal surface in dimensions  $2m \geq 14$  for all  $s \in (0, 1/2)$ .

First analytical proof of a stability result for the Simons cone in any dimension (in the nonlocal setting).

# General kernels

Rewrite the operator for doubly radial odd functions in the form:

$$L_K w(x) = \int_{\mathcal{O}} \{w(x) - w(y)\} \{\bar{K}(x, y) - \bar{K}(x, y^*)\} dy + 2w(x) \int_{\mathcal{O}} \bar{K}(x, y^*) dy,$$

where  $\bar{K}$  is doubly radial in both variables and is defined by

$$\bar{K}(x, y) := \int_{\mathcal{O}(m)^2} K(|Rx - y|) dR.$$

We establish a **necessary and sufficient condition** on the kernel  $K$  such that a theory of existence and uniqueness of saddle-shaped solutions can be developed.

This condition turns out to be

$K(\sqrt{\tau})$  is a **strictly convex** function of  $\tau$ .

# New results

		Local case	Nonlocal	
			Fractional Laplacian	General Kernels
Existence	$2m = 2$	[Dang, Fife, Peletier, '92]	[Cinti, '11, '17]	[F-N, Sanz-Perela, '19]
	$2m \geq 4$	[Cabr�, Terra, '09]		
Asymptotic behavior	$2m = 2$	[Dang, Fife, Peletier, '92]	[Cinti, '11, '17]	[F-N, Sanz-Perela, '19]
	$2m \geq 4$	[Cabr�, Terra, '10]		
Uniqueness	$2m = 2$	[Dang, Fife, Peletier, '92]	[F-N, Sanz-Perela, '18]	[F-N, Sanz-Perela, '19]
	$2m \geq 4$	[Cabr�, '12]		
Monotonicity properties	$2m = 2$	[Dang, Fife, Peletier, '92]	[Cinti, '11, '17] [F-N, Sanz-Perela, '18]	
	$2m \geq 4$	[Cabr�, Terra, '10] [Cabr�, '12]		
Unstability	$2m = 2$	[Shatzman, '95]	[Cabr�, Sol�-Morales, '05]	
	$2m = 4, 6$	[Cabr�, Terra, '09, '10]	[Cinti, '11, '17]	
Stability	$2m \geq 14$	[Cabr�, '12]	[F-N, Sanz-Perela, '18]	
Minimality	$2m \geq 8$	?	?	




# New results: key ingredients

To establish the previous results for the saddle-shaped solution, we need to prove the following results:

- An energy estimate for doubly radial minimizers.
- A Liouville type result for nonnegative bounded solutions to  $L_K u = f(u)$  in  $\mathbb{R}^n$ .
- A one-dimensional symmetry result for positive solutions to  $L_K u = f(u)$  in a half-space  $\mathbb{R}^n \cap \{x_n > 0\}$ .
- A maximum principle in “narrow” sets.

## Conclusion: Importance of saddle-shaped solutions

- They could be used to construct a counterexample for the nonlocal analogue of the conjecture by De Giorgi on the one-dimensional symmetry of monotone solutions.
- They are expected to be, in high dimensions, the simplest example of minimizer to  $L_K u = f(u)$  which is not one-dimensional (and not monotone in one direction).
- Saddle-shaped solutions can be used to prove the stability or minimality of the Simons cone as a nonlocal minimal surface.

-  J.C. Felipe-Navarro, T. Sanz-Perela, *Uniqueness and stability of the saddle-shaped solution to the fractional Allen-Cahn equation*, preprint arXiv:1810.08483 .
-  J.C. Felipe-Navarro, T. Sanz-Perela, *Semilinear integro-differential equations, I: odd solutions with respect to the Simons cone*, preprint arXiv:1903.05158.
-  J.C. Felipe-Navarro, T. Sanz-Perela, *Semilinear integro-differential equations, II: one-dimensional and saddle-shaped solutions to the Allen-Cahn equation*, in preparation.

Thank You